

# Hyperbolic analogue of the Eastwood-Norbury formula for Atiyah determinant

Dragutin Svrtan

[dragutin.svrtan@gmail.com](mailto:dragutin.svrtan@gmail.com)

University of Zagreb

September 22, 2022

# Outline

- 1 Introduction
- 2 3 points inside circle
- 3 Normalized determinant  $D_3$
- 4 Explicit formulas for  $D_3$
- 5 4 points inside a ball
- 6 Normalized 4–point Atiyah’s determinant  $D_4$
- 7 Conjectures ( $n = 4$ )
- 8 New proof of the Eastwood–Norbury formula
- 9 Positive parametrization of distances between 4 points
- 10 Verification of the Atiyah–Sutcliffe four–point conjectures
- 11 Mixed Atiyah determinants
- 12 Main Theorem
- 13 New developments (1/3)
- 14 New developments (2/3)
- 15 New developments (3/3)
- 16 References

# Introduction 1/3

In 2001, Sir Michael Atiyah, inspired by physics (Berry–Robbins problem related to spin statistics theorem of quantum mechanics) associated a remarkable determinant to any  $n$  distinct points in Euclidean 3–space, via elementary construction. More generally, let  $(x_1, x_2, \dots, x_n)$  be  $n$  distinct points inside the ball of radius  $R$  in Euclidean 3–space.

## Definition.

Let the oriented line  $x_i x_j$  meet the boundary 2–sphere in a point (direction)  $u_{ij}$  regarded as a point of the complex Riemann sphere  $(\mathbb{C} \cup \{\infty\})$ .

Form a complex polynomial  $p_i$  of degree  $n - 1$  whose roots are  $u_{ij}, j \neq i$  ( $p_i$  is determined up to a scalar factor). The Atiyah's conjecture  $C_1$  now reads

## Conjecture $C_1$

For all  $(x_1, x_2, \dots, x_n)$  the  $n$  polynomials  $p_i$  are linearly independent.

Conjecture  $C_1 \Leftrightarrow$  nonvanishing of the determinant  $D$  of the matrix of coefficients of the polynomials  $p_i$ .

The determinant  $D$  can be normalized so that  $D$  becomes a continuous function of  $(x_1, x_2, \dots, x_n)$  which is  $SL(2, \mathbb{C})$ –invariant (using the ball model or upper half space model of hyperbolic 3–space).

## Introduction 2/3

The more refined conjectures of Atiyah and Sutcliffe  $C_2$  and  $C_3$  relate  $D$  to products of 2 and  $n - 1$ -subsequences of points  $x_1, x_2, \dots, x_n$ .

The conjecture  $C_1$  is proved for  $n = 3, 4$  and for general  $n$  only for some special configurations (M.F. Atiyah, M. Eastwood and P. Norbury, D. Đoković).

In a lengthy preprint [5] we have verified the conjectures  $C_2$  and  $C_3$  for parallelograms, cyclic quadrilaterals and some infinite families of tetrahedra.

Also we proved  $C_2$  for Đoković's dihedral configurations. In [8] a proof of  $C_1$  is given for convex planar quadrilaterals. We have also proposed a strengthening of the conjecture  $C_3$  for configurations of four points (Four Points Conjecture, stronger than some new conjectures in [8]) and a number of conjectures for almost collinear configurations, and proved them for  $n$  up to 10.

In 2001, Eastwood and Norbury [3] found an intrinsic formula for the four point Atiyah determinant (a polynomial of sixth degree in six distances having several hundreds of terms) and gave a proof of  $C_1$ .

## Introduction 3/3


The present author found a new geometric fact for arbitrary tetrahedra which leads to a proof of  $C_2$  and  $C_3$  for arbitrary four points in the euclidean three space (and also a proof of stronger Four Points Conjecture of Svrtan and Urbih).

Later we obtain another intrinsic polynomial formula a la Eastwood and Norbury for four points (and for five "planar" points – having one hundred thousand terms) and have an existence proof of a polynomial formula for all planar configurations what was conjectured in [3].

This approach produces also trigonometric formulas for four points Atiyah determinants (not involving so called Crell angles which are used in [8]). Some work is done in the hyperbolic case by finding a hyperbolic analogue of the Eastwood and Norbury formula (in the planar case- spacial case is quite a challenge!).

We also introduce (mixed) Atiyah type energies associated to any graph (on given points) and can prove that Conjecture  $C_1$  is true, for arbitrary  $n$ , for some of these energies (work in progress).

# 3 points inside circle



- Three points  $x_1, x_2, x_3$  inside disk ( $|z| \leq R$ )
- Three point-pairs on circle
- $P_1 \quad (u_{12})(u_{13})$
- $P_2 \quad (u_{21})(u_{23})$
- $P_3 \quad (u_{31})(u_{32})$
- Point-pair  $u_{12}, u_{13}$  define quadratic with roots  
$$p_1 = (z - u_{12})(z - u_{13})$$
- 3 point-pairs  $\rightarrow$  3 quadratics
- $P_1, P_2, P_3 \rightarrow \{p_1, p_2, p_3\}$

## Theorem (Atiyah 2001.)

For any triple  $x_1, x_2, x_3$  of distinct points inside the disk the three quadratics  $\{p_1, p_2, p_3\}$  are linearly independent.

Remark: Atiyah's proof, which is synthetic, does not generalize to more than three points.

# Normalized determinant $D_3$

## Theorem 1.

3-by-3 determinant of the coefficient matrix:

$$|M_3| = \begin{vmatrix} 1 & -u_{12} - u_{13} & u_{12}u_{13} \\ 1 & -u_{21} - u_{23} & u_{21}u_{23} \\ 1 & -u_{31} - u_{32} & u_{31}u_{32} \end{vmatrix} \neq 0, \quad D_3 = \frac{|M_3|}{(u_{12}-u_{21})(u_{13}-u_{31})(u_{23}-u_{32})}$$

Remark:  $D_3 = 1$  only for collinear points.

## Theorem 2.

$$D_3 \geq 1.$$

Remark: Theorem 2.  $\Leftrightarrow$  Theorem 1.

Points on the "circle at  $\infty$ " are directions on a plane.

Remark: Theorem 1. and Theorem 2. are also true for  $R = \infty$ .

## Explicit formulas for $D_3$

Extrinsic formula:

$$D_3 = 1 + \frac{(u_{21} - u_{31})(u_{13} - u_{23})(u_{12} - u_{32})}{(u_{12} - u_{21})(u_{13} - u_{31})(u_{23} - u_{32})}$$

Intrinsic formula for hyperbolic triangles ( $0 < A + B + C < \pi$ ):

$$D_3 = \frac{1}{2}(\cos^2(A/2) + \cos^2(B/2) + \cos^2(C/2)) - \frac{1}{4}\Phi,$$

where  $\Phi^2 = 4 \cos\left(\frac{A+B+C}{2}\right) \cos\left(\frac{-A+B+C}{2}\right) \cos\left(\frac{A-B+C}{2}\right) \cos\left(\frac{A+B-C}{2}\right)$   
 $= -1 + \cos^2(A) + \cos^2(B) + \cos^2(C) + 2 \cos(A) \cos(B) \cos(C)$

Intrinsic formula involving side lengths

$$a, b, c, p = (a + b + c)/2, p_a = p - a, p_b = p - b, p_c = p - c:$$

$$\begin{aligned} D_3 &= 1 + e^{-p} \frac{\sinh(p_a) \sinh(p_b) \sinh(p_c)}{\sinh(a) \sinh(b) \sinh(c)} \left( \rightarrow 1 + \frac{(-a+b+c)(a-b+c)(a+b-c)}{8abc} \text{ Eucl. case} \right) \\ &= 1 + e^{-(p_a+p_b+p_c)} \frac{(e^{p_a} - e^{-p_a})(e^{p_b} - e^{-p_b})(e^{p_c} - e^{-p_c})}{(e^{p_a+p_b} - e^{-(p_a+p_b)})(e^{p_a+p_c} - e^{-(p_a+p_c)})(e^{p_b+p_c} - e^{-(p_b+p_c)})} \\ &= 1 + \frac{(e^{2p_a} - 1)(e^{2p_b} - 1)(e^{2p_c} - 1)}{(e^{2(p_a+p_b)} - 1)(e^{2(p_a+p_c)} - 1)(e^{2(p_b+p_c)} - 1)} \end{aligned}$$

$$D_3 = 1 + \frac{(e^{2p_a} - 1)(e^{2p_b} - 1)(e^{2p_c} - 1)}{(e^{2(p_a+p_b)} - 1)(e^{2(p_a+p_c)} - 1)(e^{2(p_b+p_c)} - 1)}$$

## Lemma.

For  $0 < a < b$  the function  $f(x) = \frac{e^{\frac{a}{x}} - 1}{e^{\frac{b}{x}} - 1}$  ( $0 < x < \infty$ ) is strictly increasing and  $\lim_{x \rightarrow \infty} f(x) = \frac{a}{b}$ .

By using this lemma the recent **monotonicity conjecture** of Atiyah (in case  $n = 3$ ) follows immediately (if  $a$  is replaced by  $a/R$  etc... in previous formulas).

$$\begin{aligned} D_3 &= 1 + e^{-p} \frac{\sinh(p_a) \sinh(p_b) \sinh(p_c)}{\sinh(a) \sinh(b) \sinh(c)} = 1 + \frac{e^{-p_a-p_b-p_c} \sinh(p_a) \sinh(p_b) \sinh(p_c)}{\sinh(p_a + p_b) \sinh(p_a + p_c) \sinh(p_b + p_c)} \\ &= 1 + \frac{(\cosh(p_a + p_b + p_c) - \sinh(p_a + p_b + p_c)) \sinh(p_a) \sinh(p_b) \sinh(p_c)}{\sinh(p_a + p_b) \sinh(p_a + p_c) \sinh(p_b + p_c)} \\ &= 1 + \frac{(1 - \tanh(p_a))(1 - \tanh(p_b))(1 - \tanh(p_c)) \tanh(p_a) \tanh(p_b) \tanh(p_c)}{(\tanh(p_a) + \tanh(p_b))(\tanh(p_a) + \tanh(p_c))(\tanh(p_b) + \tanh(p_c))} \end{aligned}$$

# Equations for Atiyah 3pt energies (1/4)

$$\begin{aligned}\Delta_3 &= D_3 - 1 = e^{-p} \frac{\sinh(p-a)\sinh(p-b)\sinh(p-c)}{\sinh(a)\sinh(b)\sinh(c)}, \quad \left( p = \frac{a+b+c}{2} \right) \\ \Delta_3^+ &= D_3^+ - 1 = e^p \frac{\sinh(p-a)\sinh(p-b)\sinh(p-c)}{\sinh(a)\sinh(b)\sinh(c)} \\ P_1 &= (X - \Delta_3)(X - \Delta_3^+) = \\ &= X^2 - 2 \frac{\cosh(p)\sinh(p-a)\sinh(p-b)\sinh(p-c)}{\sinh(a)\sinh(b)\sinh(c)} X + \frac{\sinh^2(p-a)\sinh^2(p-b)\sinh^2(p-c)}{\sinh^2(a)\sinh^2(b)\sinh^2(c)} \\ \Delta_3^{(1)} &= D_3^{001} - 1 = -e^{-(p-c)} \frac{\sinh(p)\sinh(p-a)\sinh(p-b)}{\sinh(a)\sinh(b)\sinh(c)} = -e^{-p+c} \frac{\sinh(p)}{\sinh(c)} \sin^2(A) \\ \Delta_3^{(6)} &= D_3^{110} - 1 = -e^{p-c} \frac{\sinh(p)\sinh(p-a)\sinh(p-b)}{\sinh(a)\sinh(b)\sinh(c)} = -e^{+p-c} \frac{\sinh(p)}{\sinh(c)} \sin^2(A) \\ P_2 &= (X - \Delta_3^{(1)})(X - \Delta_3^{(6)}) = \\ &= X^2 - 2 \frac{\sinh(p)\sinh(p-a)\sinh(p-b)\cosh(p-c)}{\sinh(a)\sinh(b)\sinh(c)} X + \frac{\sinh^2(p)\sinh^2(p-a)\sinh^2(p-b)}{\sinh^2(a)\sinh^2(b)\sinh^2(c)} \\ P_3 &= (X - \Delta_3^{(2)})(X - \Delta_3^{(5)}) = \\ &= X^2 - 2 \frac{\sinh(p)\sinh(p-a)\cosh(p-b)\sinh(p-c)}{\sinh(a)\sinh(b)\sinh(c)} X + \frac{\sinh^2(p)\sinh^2(p-a)\sinh^2(p-c)}{\sinh^2(a)\sinh^2(b)\sinh^2(c)} \\ P_4 &= (X - \Delta_3^{(3)})(X - \Delta_3^{(4)}) = \\ &= X^2 - 2 \frac{\sinh(p)\cosh(p-a)\sinh(p-b)\sinh(p-c)}{\sinh(a)\sinh(b)\sinh(c)} X + \frac{\sinh^2(p)\sinh^2(p-b)\sinh^2(p-c)}{\sinh^2(a)\sinh^2(b)\sinh^2(c)}\end{aligned}$$

# Equations for Atiyah 3pt energies (2/4)

$$\begin{aligned} P(X) &= P_1 P_2 P_3 P_4 = \prod_{i=0}^7 (X - \Delta^{(i)}) = X^8 + \cdots + e_7 X - e_8 \\ e_8 &= \prod_{i=0}^7 \Delta^{(i)} = \frac{\sinh^6(p) \sinh^6(p-a) \sinh^6(p-b) \sinh^6(p-c)}{\sinh^6(a) \sinh^6(b) \sinh^6(c)} \\ e_8 &= \frac{(1-c_1^2)^2 (1-c_2^2)^2 (1-c_3^2)^2}{4096} = \frac{\sin^4(A) \sin^4(B) \sin^4(C)}{4096} \\ e_7 &= \frac{1}{256} (1-c_1^2)(1-c_2^2)(1-c_3^2)(-1+2c_1c_2c_3+c_1^2+c_2^2+c_3^2) = \frac{1}{256} \sin^2(A) \sin^2(B) \sin^2(C) \Phi^2 \\ P(X) &= X^8 + 2X^7 + \frac{1}{4}(4+\sigma_1-\Phi^2)X^6 + \frac{1}{4}(\sigma_1-2\Phi^2)X^5 - \frac{1}{32}(3\sigma_3-2\sigma_2+(2\sigma_1-8)\Phi^2)X^4 - \\ &\quad - \frac{1}{32}(\sigma_3+2\sigma_1\Phi^2)X^3 - \frac{1}{256}(\sigma_1\sigma_3+(4\sigma_2-\sigma_3)\Phi^2)X^2 - \frac{1}{256}\sigma_3\Phi^2X + \frac{1}{4096}\sigma_3^2 \\ &\text{where } \sigma_1 = \sin^2(A) + \sin^2(B) + \sin^2(C), \sigma_2 = \sin^2(A) \sin^2(B) + \sin^2(A) \sin^2(C) + \sin^2(B) \sin^2(C), \\ &\quad \sigma_3 = \sin^2(A) \sin^2(B) \sin^2(C), \Phi^2 = 4 \cos(\sigma) \cos(\sigma-A) \cos(\sigma-B) \cos(\sigma-C), \sigma = \frac{A+B+C}{2} \end{aligned}$$

$$\prod_{i=0}^7 D_3^{(i)} = \frac{1}{128}(\sigma_3-2\sigma_2)\Phi^2 + \frac{1}{4096}(\sigma_3^2-16\sigma_1\sigma_3+256(\sigma_2-\sigma_3))$$

# Equations for Atiyah 3pt energies (3/4)

$$\begin{aligned}P_-(X) &= X^4 + (1 + \Phi)X^3 + \frac{1}{4}(c_1^2 + c_2^2 + c_3^2 + 3c_1c_2c_3 + 3\Phi)X^2 + \frac{1}{8}(-1 + 2c_1c_2c_3 + c_1^2 + c_2^2 + c_3^2 + \\&\quad + (1 + c_1c_2c_3)\Phi)X - \frac{1}{64}(1 - c_1^2)(1 - c_2^2)(1 - c_3^2) \\&= \frac{1}{8}X(2X + \Phi)[(2X + 1)(2X + 1 + \Phi) + c_1c_2c_3] - \frac{1}{64}(1 - c_1^2)(1 - c_2^2)(1 - c_3^2) \\&= \frac{1}{8}X(2X + \Phi)\left[(2X + 1)(2X + 1 + \Phi) + \frac{1 + \Phi^2 - (c_1^2 + c_2^2 + c_3^2)}{2}\right] - \frac{1}{64}(1 - c_1^2)(1 - c_2^2)(1 - c_3^2) \\&= \frac{1}{16}X(2X + \Phi)[2(2X + 1)(2X + 1 + \Phi) - 2 + s_1^2 + s_2^2 + s_3^2] - \frac{1}{64}s_1^2s_2^2s_3^2 \\P(X) &= P_-(X)P_+(X), \text{ where } P_+(X) = P_-(X) \Big|_{\Phi \rightarrow -\Phi}\end{aligned}$$

# Equations for Atiyah 3pt energies (4/4)

$$\begin{aligned} P_-(X) \Big|_{\Phi=0} &= \frac{1}{16} X \cdot 2X \left[ 2(2X+1)^2 - 2 + s_1^2 + s_2^2 + s_3^2 \right] - \frac{1}{64} s_1^2 s_2^2 s_3^2 \\ &= X^3(X+1) + \frac{1}{8}(s_1^2 + s_2^2 + s_3^2)X^2 - \frac{1}{64} s_1^2 s_2^2 s_3^2 \\ P(0) &= \frac{1}{4096} s_1^4 s_2^4 s_3^4 \\ \prod_{i=0}^7 D_3^{(i)} &= P(-1) = \left[ \frac{1}{16} (-1)(-2+\Phi)[-2(-1+\Phi) - 2 + s_1^2 + s_2^2 + s_3^2] - \frac{1}{64} s_1^2 s_2^2 s_3^2 \right] \cdot \\ &\quad \cdot \left[ \frac{1}{16} (-1)(-2-\Phi)[-2(-1-\Phi) - 2 + s_1^2 + s_2^2 + s_3^2] - \frac{1}{64} s_1^2 s_2^2 s_3^2 \right] \\ &= \left[ \frac{1}{16} (2-\Phi)[-2\Phi + \sigma_1] - \frac{1}{64} \sigma_3 \right] \left[ \frac{1}{16} (2+\Phi)[2\Phi + \sigma_1] - \frac{1}{64} \sigma_3 \right] \\ &= \left[ \frac{1}{16} (2\Phi^2 - 4\Phi + 2\sigma_1 - \Phi\sigma_1) - \frac{1}{16} \sigma_3 \right] \left[ \frac{1}{16} (2\Phi^2 + 4\Phi + 2\sigma_1 + \Phi\sigma_1) - \frac{1}{16} \sigma_3 \right] \\ &= \left[ \frac{1}{16} (2\Phi^2 + 2\sigma_1) - \frac{1}{64} \sigma_3 - \frac{1}{16} (4 + \sigma_1)\Phi \right] \left[ \frac{1}{16} (2\Phi^2 + 2\sigma_1) - \frac{1}{64} \sigma_3 + \frac{1}{16} (4 + \sigma_1)\Phi \right] \\ &= \frac{1}{64^2} \left\{ [4(\Phi^2 + \sigma_1) - \sigma_3]^2 - 4^2 (4 + \sigma_1)^2 \Phi^2 \right\}. \end{aligned}$$

## 4 points inside a ball

- Four points  $x_1, x_2, x_3, x_4$  in a ball ( $|z| \leq R$ )
- 4 point-triples on the boundary 2–sphere
- $P_1 \quad (u_{12})(u_{13})(u_{14})$
- $P_2 \quad (u_{21})(u_{23})(u_{24})$
- $P_3 \quad (u_{31})(u_{32})(u_{34})$
- $P_4 \quad (u_{41})(u_{42})(u_{43})$
- point-triple  $u_{12}, u_{13}, u_{14}$  defines a cubic (polynomial):

$$p_1 := (z - u_{12})(z - u_{13})(z - u_{14})$$

$$p_1 = z^3 - (u_{12} + u_{13} + u_{14})z^2 + (u_{12}u_{13} + u_{12}u_{14} + u_{13}u_{14})z - u_{12}u_{13}u_{14}$$

- 4 point–triples  $\rightarrow$  4 cubics
- $P_1, P_2, P_3, P_4 \rightarrow \{p_1, p_2, p_3, p_4\}$

## Normalized 4–point Atiyah’s determinant $D_4$

Determinant of the coefficient matrix of polynomials:

$$|M_4| = \begin{vmatrix} 1 & -u_{12} - u_{13} - u_{14} & u_{12}u_{13} + u_{12}u_{14} + u_{13}u_{14} & -u_{12}u_{13}u_{14} \\ 1 & -u_{21} - u_{23} - u_{24} & u_{21}u_{23} + u_{21}u_{24} + u_{23}u_{24} & -u_{21}u_{23}u_{24} \\ 1 & -u_{31} - u_{32} - u_{34} & u_{31}u_{32} + u_{31}u_{34} + u_{32}u_{34} & -u_{31}u_{32}u_{34} \\ 1 & -u_{41} - u_{42} - u_{43} & u_{41}u_{42} + u_{41}u_{43} + u_{42}u_{43} & -u_{41}u_{42}u_{43} \end{vmatrix},$$

$$D_4 = \frac{|M_4|}{(u_{12} - u_{21})(u_{13} - u_{31})(u_{14} - u_{41})(u_{23} - u_{32})(u_{24} - u_{42})(u_{34} - u_{43})}$$

# Conjectures ( $n = 4$ )

$C_1$  (Atiyah):  $D_4 \neq 0$  ( $\Leftrightarrow p_1, p_2, p_3, p_4$  lin. indep.)

$C_2$  (Atiyah–Sutcliffe):  $|D_4| \geq 1$

$C_3$  (Atiyah–Sutcliffe):  $|D_4|^2 \geq D_3(1, 2, 3) \cdot D_3(1, 2, 4) \cdot D_3(1, 3, 4) \cdot D_3(2, 3, 4)$

# Eastwood-Norbury formulas for euclidean D4

In 2001 they proved, by tricky use of MAPLE, that for  $n = 4$  points in Eucl.

3-space

$$\begin{aligned} Re(D_4) = & \quad 64abca'b'c' \\ & -4 \cdot d3(aa', bb', cc') \\ & + SUM \\ & + 288 \cdot VOLUME^2, \end{aligned}$$


where

$$SUM := a'[(b' + c')^2 - a^2]d3(a, b, c) + \dots$$

$$D_4/(64abca'b'c') = D_4$$

(=>eucl. Conjecture 1, and "almost"

(= 60/64) of euclidean Conjecture 2)



$$a'((b' + c')^2 - a^2)^* d3(a, b, c)$$

# New proof of the Eastwood–Norbury formula

The four points:

$$P_i : x_i = (z_i, r_i), z_i \in \mathbb{C}, r_i \in \mathbb{R}$$

$$R_{ij} := r_{ij} + r_i - r_j, z_{ij} := z_i - z_j$$


$$R_{ij} R_{ji} = r_{ij}^2 - (r_i - r_j)^2 = |z_{ij}|^2 = -z_{ij} z_{ji}$$

$$p_1 = \left( z + \frac{\overline{z_{12}}}{R_{12}} \right) \left( z + \frac{\overline{z_{13}}}{R_{13}} \right) \left( z + \frac{\overline{z_{14}}}{R_{14}} \right)$$

$$p_2 = \left( z + \frac{\overline{z_{21}}}{R_{21}} \right) \left( z + \frac{\overline{z_{23}}}{R_{23}} \right) \left( z + \frac{\overline{z_{24}}}{R_{24}} \right)$$

$$p_3 = \left( z + \frac{\overline{z_{31}}}{R_{31}} \right) \left( z + \frac{\overline{z_{32}}}{R_{32}} \right) \left( z + \frac{\overline{z_{34}}}{R_{34}} \right)$$

$$p_4 = \left( z + \frac{\overline{z_{41}}}{R_{41}} \right) \left( z + \frac{\overline{z_{42}}}{R_{42}} \right) \left( z + \frac{\overline{z_{43}}}{R_{43}} \right)$$



## Matrix of coefficients of $\{p_1, p_2, p_3, p_4\}$

$$M_4 = \begin{vmatrix} 1 & \cdot & \cdot & \cdot \\ 1 & \boxed{\frac{\bar{z}_{21}}{R_{21}} + \frac{\bar{z}_{23}}{R_{23}} + \frac{\bar{z}_{24}}{R_{24}}} & \cdot & \frac{\bar{z}_{12}}{R_{12}} \frac{\bar{z}_{13}}{R_{13}} \frac{\bar{z}_{14}}{R_{14}} \\ 1 & \cdot & \boxed{\frac{\bar{z}_{31}}{R_{31}} \frac{\bar{z}_{32}}{R_{32}} + \frac{\bar{z}_{31}}{R_{31}} \frac{\bar{z}_{34}}{R_{34}} + \frac{\bar{z}_{32}}{R_{32}} \frac{\bar{z}_{34}}{R_{34}}} & \cdot \\ 1 & \cdot & \cdot & \boxed{\frac{\bar{z}_{41}}{R_{41}} \frac{\bar{z}_{42}}{R_{42}} \frac{\bar{z}_{43}}{R_{43}}} \\ \end{vmatrix} \cdot A \\ \cdot B \\ \cdot C$$

$A = z_{21}, B = z_{31}z_{32}, C = z_{41}z_{42}z_{43}$

## Normalized Atiyah determinant

$$\begin{aligned}
 D_4 &= \underbrace{\det(M_4)}_{antisym.} \cdot \underbrace{z_{21} \cdot z_{31}z_{32} \cdot z_{41}z_{42}z_{43}}_{antisym.} = \sum 1 \cdot \left( \frac{z_{21}\bar{z}_{21}}{R_{21}} + \frac{z_{21}\bar{z}_{23}}{R_{23}} + \frac{z_{21}\bar{z}_{24}}{R_{24}} \right) \cdot \\
 &\quad \cdot \left( \frac{z_{31}\bar{z}_{31}z_{32}\bar{z}_{32}}{R_{31}R_{32}} + \frac{z_{31}\bar{z}_{31}z_{32}\bar{z}_{34}}{R_{31}R_{34}} + \frac{z_{32}\bar{z}_{32}z_{32}\bar{z}_{34}}{R_{32}R_{34}} \right) R_{14}R_{24}R_{34} = \\
 &= \sum (R_{12}R_{24} + \underbrace{z_{21}z_{24}}_{}) (R_{13}R_{23}R_{34} + R_{13} \underbrace{z_{32}\bar{z}_{34}}_{} + R_{23} \underbrace{z_{31}\bar{z}_{34}}_{}) R_{14} + \\
 &+ (R_{13}R_{24}R_{34} \underbrace{z_{21}\bar{z}_{23}}_{} + R_{13}R_{24}R_{32} \underbrace{z_{21}\bar{z}_{34}}_{} + R_{24} \underbrace{z_{21}\bar{z}_{23}z_{31}\bar{z}_{34}}_{}) R_{14}
 \end{aligned}$$

(where summations are over all permutations of indices).

By writing  $z_{ij}\bar{z}_{kl} = C[i, j, k, l] + \sqrt{-1} S[i, j, k, l]$  and using a Lagrange identity (involving the dot product of two cross products; a fact mentioned by N. Wildberger to the author) we have

$S[i, j, k, l]S[p, q, r, s] = C[i, j, p, q]C[k, l, r, s] - C[i, j, r, s]C[k, l, p, q]$   
(we have discovered this identity independently) and using the formula

$$\begin{aligned} C[i, j, k, l] &= \operatorname{Re}(z_{ij}\bar{z}_{kl}) = \frac{1}{2}[|z_{il}|^2 + |z_{jk}|^2 - |z_{ik}|^2 - |z_{jl}|^2] = \\ &= \frac{1}{2}[r_{il}^2 + r_{jk}^2 - r_{ik}^2 - r_{jl}^2] - (r_i - r_j)(r_k - r_l) \end{aligned}$$

we obtain our derivation of the Eastwood–Norbury formula.

By this new method we obtained a polynomial formula for the planar configurations of 5 points (by  $S_5$ –symmetrization of a "one page" expression) and a rational formula for the spatial 5 point configuration (this last formula has almost 100000 terms).

This settles one of the Eastwood–Norbury conjectures. We do not yet have definite geometric interpretations for the "nonplanar" part of the formula involving heights  $r_i$ ,  $i = 1, \dots, 5$ .



Our trigonometric (euclidean) Eastwood–Norbury formula  
(where  $c_{i\_jk} := \cos(ij, ik)$  and  $c_{ij\_kl} := \cos(ij, kl)$  ):

$$\begin{aligned} 16Re(D_4) = & (1 + c_{3\_12} + c_{2\_34})(1 + c_{1\_24} + c_{4\_13}) + \\ & (1 + c_{2\_13} + c_{3\_24})(1 + c_{4\_12} + c_{1\_34}) + \\ & (1 + c_{3\_12} + c_{1\_34})(1 + c_{2\_14} + c_{4\_23}) + \\ & (1 + c_{1\_23} + c_{3\_14})(1 + c_{2\_34} + c_{4\_12}) + \\ & (1 + c_{2\_13} + c_{1\_24})(1 + c_{3\_14} + c_{4\_23}) + \\ & (1 + c_{1\_23} + c_{2\_14})(1 + c_{3\_24} + c_{4\_13}) + \\ & 2(c_{14\_23}c_{13\_24} - c_{14\_23}c_{12\_34} + c_{13\_24}c_{12\_34}) + \\ & 72(\text{normalized volume})^2. \end{aligned}$$

Open problems:

Hyperbolic (euclidean) version for  $n \geq 4$  ( $n \geq 5$ ) points in terms of  
distances, or in terms of angles.

# NEW TRIGONOMETRIC FORMULA FOR NORMALIZED VOLUME OF A TETRAHEDRON

A law of sines for tetrahedra and the space of all shapes of tetrahedra

A corollary of the usual law of sines is that in a tetrahedron with vertices O, A, B, C, we have

$$\sin \angle OAB \cdot \sin \angle OBC \cdot \sin \angle OCA = \sin \angle OAC \cdot \sin \angle OCB \cdot \sin \angle OBA.$$

Putting any of the four vertices in the role of O yields four such identities, but at most three of them are independent: If the "clockwise" sides of three of them are multiplied and the product is inferred to be equal to the product of the "counterclockwise" sides of the same three identities, and then common factors are canceled from both sides, the result is the fourth identity. Three angles are the angles of some triangle if and only if their sum is  $180^\circ$  ( $\pi$  radians). What condition on 12 angles is necessary and sufficient for them to be the 12 angles of some tetrahedron?

Clearly the sum of the angles of any side of the tetrahedron must be  $180^\circ$ . Since there are four such triangles, there are four such constraints on sums of angles, and the number of degrees of freedom is thereby reduced from 12 to 8. The four relations given by this sine law further reduce the number of degrees of freedom, from 8 down to not 4 but 5, since the fourth constraint is not independent of the first three. Thus the space of all shapes of tetrahedra is 5-dimensional.

Recall the Cayley-Menger matrix, the squared volume of a tetrahedron:

$$M_4 = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & r_{12}^2 & r_{13}^2 & r_{14}^2 \\ 1 & r_{12}^2 & 0 & r_{23}^2 & r_{24}^2 \\ 1 & r_{13}^2 & r_{23}^2 & 0 & r_{34}^2 \\ 1 & r_{14}^2 & r_{24}^2 & r_{34}^2 & 0 \end{bmatrix}$$

$$Vsq := \frac{\text{Determinant}(M_4)}{2^3 (3!)^2}$$

$$\begin{aligned}
Vsq := & - \frac{r_{12}^4 r_{34}^2}{144} - \frac{r_{12}^2 r_{13}^2 r_{23}^2}{144} + \frac{r_{12}^2 r_{13}^2 r_{24}^2}{144} + \frac{r_{12}^2 r_{13}^2 r_{34}^2}{144} \\
& + \frac{r_{12}^2 r_{14}^2 r_{23}^2}{144} - \frac{r_{12}^2 r_{14}^2 r_{24}^2}{144} + \frac{r_{12}^2 r_{14}^2 r_{34}^2}{144} + \frac{r_{12}^2 r_{23}^2 r_{34}^2}{144} \\
& + \frac{r_{12}^2 r_{24}^2 r_{34}^2}{144} - \frac{r_{12}^2 r_{34}^4}{144} - \frac{r_{13}^4 r_{24}^2}{144} + \frac{r_{13}^2 r_{14}^2 r_{23}^2}{144} \\
& + \frac{r_{13}^2 r_{14}^2 r_{24}^2}{144} - \frac{r_{13}^2 r_{14}^2 r_{34}^2}{144} + \frac{r_{13}^2 r_{23}^2 r_{24}^2}{144} - \frac{r_{13}^2 r_{24}^4}{144} \\
& + \frac{r_{13}^2 r_{24}^2 r_{34}^2}{144} - \frac{r_{14}^4 r_{23}^2}{144} - \frac{r_{14}^2 r_{23}^4}{144} + \frac{r_{14}^2 r_{23}^2 r_{24}^2}{144} \\
& + \frac{r_{14}^2 r_{23}^2 r_{34}^2}{144} - \frac{r_{23}^2 r_{24}^2 r_{34}^2}{144}
\end{aligned}$$

$$vols1 := 144Vsq$$

$$\begin{aligned} vols1 := & -r_{12}^4 r_{34}^2 - r_{12}^2 r_{13}^2 r_{23}^2 + r_{12}^2 r_{13}^2 r_{24}^2 + r_{12}^2 r_{13}^2 r_{34}^2 \\ & + r_{12}^2 r_{14}^2 r_{23}^2 - r_{12}^2 r_{14}^2 r_{24}^2 + r_{12}^2 r_{14}^2 r_{34}^2 + r_{12}^2 r_{23}^2 r_{34}^2 \\ & + r_{12}^2 r_{24}^2 r_{34}^2 - r_{12}^2 r_{34}^4 - r_{13}^4 r_{24}^2 + r_{13}^2 r_{14}^2 r_{23}^2 \\ & + r_{13}^2 r_{14}^2 r_{24}^2 - r_{13}^2 r_{14}^2 r_{34}^2 + r_{13}^2 r_{23}^2 r_{24}^2 - r_{13}^2 r_{24}^4 \\ & + r_{13}^2 r_{24}^2 r_{34}^2 - r_{14}^4 r_{23}^2 - r_{14}^2 r_{23}^4 + r_{14}^2 r_{23}^2 r_{24}^2 \\ & + r_{14}^2 r_{23}^2 r_{34}^2 - r_{23}^2 r_{24}^2 r_{34}^2 \end{aligned}$$

To each vertex  $i = 1..4$  and a cyclic orientation of its complement we associate the following quantities:

$$b1 := (r_{12}^2 - r_{13}^2 + r_{23}^2)(r_{13}^2 - r_{14}^2 + r_{34}^2)(-r_{12}^2 + r_{14}^2 + r_{24}^2) + \\ + (-r_{12}^2 + r_{13}^2 + r_{23}^2)(-r_{13}^2 + r_{14}^2 + r_{34}^2)(r_{12}^2 - r_{14}^2 + r_{24}^2)$$

$$b2 := (r_{12}^2 + r_{13}^2 - r_{23}^2)(r_{23}^2 - r_{24}^2 + r_{34}^2)(-r_{12}^2 + r_{14}^2 + r_{24}^2) + \\ + (-r_{12}^2 + r_{13}^2 + r_{23}^2)(-r_{23}^2 + r_{24}^2 + r_{34}^2)(r_{12}^2 + r_{14}^2 - r_{24}^2)$$

$$b3 := (r_{12}^2 + r_{13}^2 - r_{23}^2)(r_{23}^2 + r_{24}^2 - r_{34}^2)(-r_{13}^2 + r_{14}^2 + r_{34}^2) + \\ + (r_{12}^2 - r_{13}^2 + r_{23}^2)(-r_{23}^2 + r_{24}^2 + r_{34}^2)(r_{13}^2 + r_{14}^2 - r_{34}^2)$$

$$b4 := (r_{12}^2 + r_{14}^2 - r_{24}^2)(r_{23}^2 + r_{24}^2 - r_{34}^2)(r_{13}^2 - r_{14}^2 + r_{34}^2) + \\ + (r_{12}^2 - r_{14}^2 + r_{24}^2)(r_{23}^2 - r_{24}^2 + r_{34}^2)(r_{13}^2 + r_{14}^2 - r_{34}^2)$$

Then it follows that

$$4 \text{ vols1} = b_1 + b_2 + b_3 + b_4.$$

Recall the notation for cosine of angle at vertex  $i$  in a triangle with vertices  $i, j, k$ :

$$C_{i\_jk} = \frac{r_{ij}^2 + r_{ik}^2 - r_{jk}^2}{2r_{ij}r_{ik}}.$$

Then the last computation reads as the following NEW FORMULA:  
(for the normalized squared volume of a tetrahedron)

$$\frac{288V^2}{64 r_{12} r_{13} r_{14} r_{23} r_{24} r_{34}} = \frac{1}{16} \sum C_{i\_jl} C_{j\_kl} C_{k\_il}$$

where summation of triple products of cosines is over all 8 oriented three – cycles  $\langle ijk \rangle$  of vertices of our tetrahedron.

**Corollary** For a semiregular tetrahedron ( $r_{12} = r_{34} = c$ ,  $r_{13} = r_{24} = b$ , and  $r_{14} = r_{23} = a$ ) we obtain the well known formula:

$$72V^2 = (-a^2 + b^2 + c^2)(a^2 - b^2 + c^2)(a^2 + b^2 - c^2).$$

# INTRINSIC FORMULA FOR THE HYPERBOLIC 4-POINT ATIYAH DETERMINANT

$$p_1 := (z - t_{12})(z - t_{13})(z - t_{14})$$

$$p_2 := (z - t_{21})(z - t_{23})(z - t_{24})$$

$$p_3 := (z - t_{31})(z - t_{32})(z - t_{34})$$

$$p_4 := (z - t_{41})(z - t_{42})(z - t_{43})$$

$$M_4 := \begin{bmatrix} 1 & -t_{12} - t_{13} - t_{14} & t_{12}t_{13} + t_{14}t_{12} + t_{14}t_{13} & -t_{12}t_{13}t_{14} \\ 1 & -t_{21} - t_{23} - t_{24} & t_{21}t_{23} + t_{24}t_{21} + t_{24}t_{23} & -t_{21}t_{23}t_{24} \\ 1 & -t_{31} - t_{32} - t_{34} & t_{31}t_{32} + t_{34}t_{31} + t_{34}t_{32} & -t_{31}t_{32}t_{34} \\ 1 & -t_{41} - t_{42} - t_{43} & t_{41}t_{42} + t_{43}t_{41} + t_{43}t_{42} & -t_{41}t_{42}t_{43} \end{bmatrix}$$

$$\Delta_4 := \text{Determinant}(M_4)$$

$$\text{prod} := (t_{12} - t_{21})(t_{13} - t_{31})(t_{23} - t_{32})(t_{14} - t_{41})(t_{24} - t_{42})(t_{34} - t_{43})$$

$$D_4 := \frac{\Delta_4}{\text{prod}} = \frac{t_{21}t_{41}t_{42}t_{43}t_{31}t_{32} + \cdots \text{ 214 more similar terms} \cdots + t_{34}t_{12}t_{13}t_{14}t_{24}t_{23}}{(t_{12} - t_{21})(t_{13} - t_{31})(t_{23} - t_{32})(t_{14} - t_{41})(t_{24} - t_{42})(t_{34} - t_{43})}$$

Let  $D_{4a}$  be equal to  $D_4$  after substitutions

$$\begin{aligned} t_{ij} &= (s A_{ij} + B_{ij})/2, i = 1..j-1, j = 1..4, \\ t_{ji} &= (s A_{ij} - B_{ij})/2, i = 1..j-1, j = 1..4. \end{aligned}$$

$$cc(i, j, k, l) := \frac{(t_{kl} - t_{ji})(t_{ij} - t_{lk})}{(t_{ij} - t_{ji})(t_{kl} - t_{lk})}$$

$$C(i, j, k, l) := 2 \frac{(t_{kl} - t_{ji})(t_{ij} - t_{lk})}{(t_{ij} - t_{ji})(t_{kl} - t_{lk})} - 1$$

## AMPLITUDE(12, 13, 24)

$$Q_{121324} := -1 + C(1,2,1,3)^2 + C(1,2,2,4)^2 + C(1,3,2,4)^2 + 2C(1,2,1,3)C(1,2,4,2)C(1,3,2,4)$$

$$Q_{121324} := \frac{4(t_{12}t_{13}t_{21} - + \cdots \text{ 10 more similar terms } \cdots + -t_{24}t_{31}t_{42})^2}{(t_{12} - t_{21})^2(t_{13} - t_{31})^2(t_{24} - t_{42})^2}$$

$Q_{121324}$  is symmetric in  $A_{ij} = t_{ij} + t_{ji}$ , and antisymmetric in  $B_{ij} = t_{ij} - t_{ji}$  coordinates:

$$\begin{aligned} Q_{121324} = & \frac{1}{4} \frac{(A_{13}-A_{24})(A_{12}-A_{24})(A_{12}-A_{13})s^3}{B_{12}B_{13}B_{24}} - \frac{1}{4} \frac{(A_{13}-A_{24})B_{12}s}{B_{13}B_{24}} \\ & + \frac{1}{4} \frac{(A_{12}-A_{24})B_{13}s}{B_{24}B_{12}} - \frac{1}{4} \frac{(A_{12}-A_{13})B_{24}s}{B_{12}B_{13}} \end{aligned}$$

$$F_{123} = \frac{1}{2} \left( \cos\left(\frac{A}{2}\right)^2 + \cos\left(\frac{B}{2}\right)^2 + \cos\left(\frac{C}{2}\right)^2 \right)$$

$$G_{123} = (-1 + \cos(A)^2 + \cos(B)^2 + \cos(C)^2 + 2 \cos(A) \cos(B) \cos(C))^{\frac{1}{2}}$$



# Positive parametrization of distances between 4 points

**Key Lemma. (Shear coordinates of a tetrahedron)**

In any tetrahedron (degenerate or not) one has the following type of nonnegative splitting of edge lengths:


$$r_{12} = t_1 + u + v + t_2, r_{13} = t_1 + v + t_3,$$

$$r_{23} = t_2 + u + t_3, r_{14} = t_1 + u + t_4,$$

$$r_{24} = t_2 + v + t_4, r_{34} = t_3 + u + v + t_4$$

if and only if  $r_{12} + r_{34} =$

$$\max\{r_{12} + r_{34}, r_{13} + r_{24}, r_{14} + r_{23}\}.$$



## Proof.

The form of the solution:

$$t_1 = \frac{r_{13} + r_{14} - r_{34}}{2}, t_2 = \frac{r_{23} + r_{24} - r_{34}}{2}, t_3 = \frac{r_{13} + r_{23} - r_{12}}{2},$$

$$t_4 = \frac{r_{14} + r_{24} - r_{12}}{2}, u = \frac{r_{12} + r_{34} - (r_{13} + r_{24})}{2}, v = \frac{r_{12} + r_{34} - (r_{14} + r_{23})}{2}$$

proves the Lemma immediately.

4


1

2

v

3

u



# Verification of the Atiyah–Sutcliffe four–point conjectures

Let us recall the original Eastwood–Norbury formula for the real part of the Atiyah’s determinant  $D_4$  of a tetrahedron:

$$Re(D_4) := prod - 4d_3(r_{12}r_{34}, r_{13}r_{24}, r_{23}r_{14}) + A_4 + vols;$$

where  $d_3(a, b, c) := (-a + b + c)(a - b + c)(a + b - c);$

$$A_4 =$$

$$\begin{aligned} & (r_{14}((r_{24} + r_{34})^2 - r_{23}^2) + r_{24}((r_{14} + r_{34})^2 - r_{13}^2) + r_{34}((r_{24} + r_{14})^2 - r_{12}^2))d_3(r_{12}, r_{13}, r_{23}) + \\ & + (r_{13}((r_{23} + r_{34})^2 - r_{24}^2) + r_{23}((r_{13} + r_{34})^2 - r_{14}^2) + r_{34}((r_{23} + r_{13})^2 - r_{12}^2))d_3(r_{12}, r_{14}, r_{24}) + \\ & + (r_{12}((r_{23} + r_{24})^2 - r_{34}^2) + r_{23}((r_{12} + r_{24})^2 - r_{14}^2) + r_{24}((r_{23} + r_{12})^2 - r_{13}^2))d_3(r_{13}, r_{14}, r_{34}) + \\ & + (r_{12}((r_{13} + r_{14})^2 - r_{34}^2) + r_{13}((r_{12} + r_{14})^2 - r_{24}^2) + r_{14}((r_{13} + r_{12})^2 - r_{23}^2))d_3(r_{23}, r_{24}, r_{34}); \end{aligned}$$

$$prod := 64r_{12}r_{13}r_{23}r_{14}r_{24}r_{34};$$

$$vols := 2(r_{12}^2r_{34}^2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2 - r_{12}^2 - r_{34}^2) + r_{13}^2r_{24}^2(-r_{13}^2 + r_{14}^2 + r_{23}^2 - r_{24}^2 + r_{12}^2 + r_{34}^2) + r_{14}^2r_{23}^2(r_{13}^2 - r_{14}^2 - r_{23}^2 + r_{24}^2 + r_{12}^2 + r_{34}^2) - r_{12}^2r_{13}^2r_{23}^2 - r_{12}^2r_{14}^2r_{24}^2 - r_{13}^2r_{14}^2r_{34}^2 - r_{23}^2r_{24}^2r_{34}^2);$$

( $vols = 288volume^2$ ) and normalized Atiyah determinant of face triangles:

$$\delta_1 := 1 + \frac{1}{8} \frac{d_3(r_{23}, r_{24}, r_{34})}{r_{23}r_{24}r_{34}}, \quad \delta_2 := 1 + \frac{1}{8} \frac{d_3(r_{13}, r_{14}, r_{34})}{r_{13}r_{14}r_{34}},$$

$$\delta_3 := 1 + \frac{1}{8} \frac{d_3(r_{12}, r_{14}, r_{24})}{r_{12}r_{14}r_{24}}, \quad \delta_4 := 1 + \frac{1}{8} \frac{d_3(r_{12}, r_{13}, r_{23})}{r_{12}r_{13}r_{23}}.$$

We first prove a stronger four–point conjecture of Svrtan – Urbija  
 (arXiv:math0609174v1 (Conjecture 2.1 (weak version)) which implies (c.f. Proposition 2.2 in loc.cit) all three four–point conjectures  $C_1, C_2, C_3$  of Atiyah – Sutcliffe).

The substitution from the Key Lemma

$$\text{Sub} := \{r_{12} = t_1 + u + v + t_2, r_{13} = t_1 + v + t_3, r_{23} = t_2 + u + t_3, \\ r_{14} = t_1 + u + t_4, r_{24} = t_2 + v + t_4, r_{34} = t_3 + u + v + t_4\};$$

in the Maple code DifferSU :=

$$\left\{ \text{coeffs} \left( \text{expand} \left( \text{subs} \left( \text{Sub}, \frac{1}{64} \text{numer} \left( \frac{\text{Re}(D_4) - 4\text{vols}}{\text{prod}} - \frac{\delta_1^2 + \delta_2^2 + \delta_3^2 + \delta_4^2}{4} \right) \right) \right) \right) \right\};$$

gives the output DifferSU = {2, 3, 4, ..., 5328, 5564, 6036} which proves the conjecture coefficientwise.

The Maple code for the strongest Atiyah – Sutcliffe conjecture DifferAS :=

$$\left\{ \text{coeffs} \left( \text{expand} \left( \text{subs} \left( \text{Sub}, \frac{1}{64} \text{numer} \left( \left( \frac{\text{Re}(D_4) - 4\text{vols}}{\text{prod}} \right)^2 - \delta_1 \delta_2 \delta_3 \delta_4 \right) \right) \right) \right) \right\};$$

gives the output DifferAS = {64, 128, 192, ..., 233472, 246720, 261888}

(coefficients of a 4512 terms inequality of degree 12 in 6 distances).

**Remark 1.** Similarly to DifferSU one can check the upper estimate with the additional coefficient equal to 37/27.

**Remark 2.** Recently we also proved Atiyah – Sutcliffe conjecture  $C_2$  directly from the following new formula:

$$Re(D_4) = 64 \prod_{1 \leq i < j \leq 4} r_{ij} + 8d_3(r_{12}r_{34}, r_{13}r_{24}, r_{14}r_{23}) + 4vols + 32R_4 ,$$

where

$$\begin{aligned} R_4 = & 4m_{2211} + (s_{13}p_{24}^2 + s_{24}p_{13}^2)u + (s_{14}p_{23}^2 + s_{23}p_{14}^2)v + (m_{221} + 8m_{2111})w + \\ & + 2(\tau_{13}^2 + \tau_{14}^2 + \tau_{13}\tau_{14})uv + (2m_{211} + 8m_{1111})(2u^2 + uv + 2v^2) \\ & + 4m_{111}(u^3 + v^3) + (3m_{21} + 14m_{111} + 3m_{11}w)uvw + [(s_{14}p_{14} + s_{23}p_{23})(u + w) + \\ & + (s_{13}p_{13} + s_{24}p_{24})(v + w)]uv + [(\tau_{13} + \tau_{14})(u^2 + uv + v^2) + \\ & + \tau_{14}u^2 + \tau_{13}v^2]uv + 2(m_1 + w)(4m_1 + 3w)u^2v^2 \end{aligned}$$

and where

$$u = \frac{r_{12} + r_{34} - r_{13} - r_{24}}{2}, \quad v = \frac{r_{12} + r_{34} - r_{14} - r_{23}}{2}, \quad w = u + v, \quad \tau_{13} = t_1t_3 + t_2t_4,$$

$$\tau_{14} = t_1t_4 + t_2t_3, \quad t_1 = \frac{r_{13} + r_{14} - r_{34}}{2}, \quad t_2 = \frac{r_{23} + r_{24} - r_{34}}{2}, \quad t_3 = \frac{r_{13} + r_{23} - r_{12}}{2},$$

$$t_4 = \frac{r_{14} + r_{24} - r_{12}}{2}, \quad s_{ij} = t_i + t_j, \quad p_{ij} = t_i t_j, \quad m_1 = t_1 + t_2 + t_3 + t_4,$$

$$m_{11} = t_1t_2 + \dots, \quad m_{21} = t_1^2t_2 + \dots, \quad m_{111} = t_1t_2t_3 + \dots, \quad m_{1111} = t_1t_2t_3t_4,$$

$$m_{2111} = t_1^2t_2t_3t_4 + \dots, \quad m_{221} = t_1^2t_2^2t_3 + \dots, \quad m_{2211} = t_1^2t_2^2t_3t_4 + \dots$$

# Mixed Atiyah determinants

We further generalize Atiyah normalized determinant  $D(x_1, \dots, x_n)$  to  $D^\Gamma(x_1, \dots, x_n)$ , where  $\Gamma$  is any (simple) graph with the vertex set  $\{x_1, \dots, x_n\}$ .

## Definition.

We start with the normalized Atiyah determinant  $D$  viewed as a function of all directions  $u_{ij}$  ( $1 \leq i \neq j \leq n$ ). Then we define  $D^\Gamma$  by simultaneously switching the roles of directions (i.e. replacing  $u_{ij}$  by  $u_{ji}$  and also replacing  $u_{ji}$  by  $u_{ij}$ ) for each pair  $ij$  such that  $x_i x_j$  is an edge of  $\Gamma$ .

For  $n = 3$  we obtain eight mixed Atiyah's determinants (mixed energies) which we can label by binary sequences  $D_3 = D_3^{000}, D_3^{001}, \dots, D_3^{111}$  for which we also have simple explicit trigonometric formulas, which can be obtained from the original Atiyah determinant by suitable sign changes of the lengths of the sides of a triangle.

Observe that

$$D_3 = D_3^{000}, D_3^{111} = 1 + e^p \prod \sinh(p_a) / \sinh(a)$$

are both  $\geq 1$ . All other mixed determinants, eg.

$$D_3^{110} = 1 - e^{p_c} \sinh(p) \sinh(p_a) \sinh(p_b) / \prod \sinh(a),$$

are between 0 and 1.

# Main Theorem

Now we state our

## Main Theorem.

We have  $\sum_{\Gamma} D^{\Gamma} = n!$ , where the summation extends over all simple graphs on  $n$  vertices.

The proof is obtained by our method of computing Atiyah's determinants.

## Corollary.

For any configuration of points in a hyperbolic 3-space at least one of the mixed Atiyah determinants is nonzero.

# Proof of the main Theorem

## Proof of the Main Theorem.

In coordinates  $B_{ij} = u_{ij} - u_{ji}$  (antisymmetric) and  $A_{ij} = u_{ij} + u_{ji}$  (symmetric)  $1 \leq i \neq j \leq n$ ,  $D^\Gamma$  differs from  $D$  in changing signs of  $B_{ij}$ 's for each edge  $ij \in \Gamma$ . Let us first observe that each nonconstant term in  $D$  (and in each  $D^\Gamma$ ) is a square free Laurent monomial w.r.t. all variables  $B_{ij}$ 's, hence in the sum over  $\Gamma$  its contribution is zero.

Therefore, we have to compute the constant term (C.T.) of  $D$  (which is the same in all  $D^\Gamma$ ). Since  $D$  is a symmetrization over  $S_n$  of its main diagonal term, we have  $C.T.(D) = n! C.T.(diagonal\ term)$ . But diagonal term of  $D$  is equal to

$$\frac{1 \cdot (-u_{21} + \dots)((-u_{31})(-u_{32}) + \dots) \cdots [(-u_{n,1})(-u_{n,1}) \cdots (-u_{n,n-1})]}{(u_{12} - u_{21})(u_{13} - u_{31})(u_{23} - u_{32}) \cdots (u_{1,n} - u_{n,1}) \cdots (u_{n-1,n} - u_{n,n-1})}$$

$$\text{so } C.T.(diag.\ term) = C.T. \frac{\frac{B_{12}}{2} \frac{B_{13}}{2} \frac{B_{23}}{2} \cdots}{B_{12} B_{13} B_{23} \cdots} = \frac{1}{2^{\binom{n}{2}}} \text{ and } C.T.(D) = \frac{n!}{2^{\binom{n}{2}}} \text{ and}$$

$$C.T. \left( \sum_{\Gamma} D^\Gamma \right) = n!. \quad \square$$

## New developments (1/3)

- In 2011, M.Mazur and B.V.Petrenko restated the original Eastwood Norbury formula in trigonometric form which besides face angles of a tetrahedron uses also angles of so called Crelle triangle (associated to the tetrahedron). Our formula in [5] does not involve Crelle's angles, but uses "skew" angles.
- $C_2$  proved for convex (planar) quadrilaterals
- $C_3$  proved for cyclic quadrilaterals (we have it proved already in [5])
- Three conjectures stated which are consequences of some of our conjectures in [5]. (Hence we have a proof of all three.)

## New developments (2/3)


- In a recent paper M.B.Khuzam and M.J.Johnson (arXiv:1401.2787v1) gave a verification (by linear programming) of both  $C_2$  and  $C_3$  four-point conjectures of Atiyah and Sutcliffe, by using symmetric functions of degree 12 in 12 variables  $t_{il} = r_{ij} + r_{ik} - r_{jk}$ ,  $\{i, j, k, l\} = \{1, 2, 3, 4\}$  (which are linearly dependent), so for  $C_2$  (resp.  $C_3$ ) they use 64 (resp. 114) huge monomial symmetric functions.
- In a recent paper J. Malkoun defined a symplectic version of Atiyah conjecture and proved it for  $n = 2$  (which also follows from [5], 2.6 Atiyah - Sutcliffe conjectures for parallelograms). We observe that symplectic Atiyah determinants are special case of ordinary Atiyah determinants for centrally symmetric configurations!

## New developments (3/3)

# Tropical Version of Atiyah–Sutcliffe Conjecture for almost collinear configurations

Almost collinear configurations. Đoković's approach and generalizations.

Type A configuration  $(n+1)$  ( $n$  points collinear)



$$\lambda_1 = a_1 + \sqrt{b^2 + a_1^2} = a_1 + d_1 = d_1(1 + \cos(\varphi_1)),$$

...

$$\lambda_2 = a_2 + \sqrt{b^2 + a_2^2} = a_2 + d_2 = d_2(1 + \cos(\varphi_2)),$$

$$\lambda_n = a_n + \sqrt{b^2 + a_n^2} = a_n + d_n = d_n(1 + \cos(\varphi_n)).$$

$$M_{n,1} = \begin{vmatrix} 1 & \lambda_1 & & & e_1 = \lambda_1 + \lambda_2 + \cdots + \lambda_n \\ & 1 & \lambda_2 & & e_2 = \lambda_1\lambda_2 + \cdots + \lambda_{n-1}\lambda_n \\ (-1)^n e_n & & -e_1 & 1 & \cdots \\ & & & & e_n = \lambda_1 \cdots \lambda_n \end{vmatrix},$$

$$\begin{aligned} D_{n,1} &= \det(M_{n,1}) = 1 + \lambda_n e_1 + \lambda_n \lambda_{n-1} e_2 + \cdots + \lambda_n \cdots \lambda_1 e_n \\ &\geq 1 + e_1(\lambda_1^2, \dots, \lambda_n^2) + \cdots + e_n(\lambda_1^2, \dots, \lambda_n^2) \\ &= (1 + \lambda_1^2)(1 + \lambda_2^2) \cdots (1 + \lambda_n^2) \Rightarrow \text{proof of } C_2 \end{aligned}$$

**Atiyah–Sutcliffe conjectures:**

$$C_2 : D_{n,1} = D_{n,1}/\prod(1 + \lambda_k^2) \geq 1$$

$$C_3 : (D_{n,1}(\lambda_1, \dots, \lambda_n))^{n-1} \geq \prod D_{n-1,1}(\lambda_1, \dots, \widehat{\lambda_k}, \dots, \lambda_n)$$

Already in 2004., we generalized asymmetric Đoković's formula for

$$D_{n,1} = 1 + \lambda_n e_1(\lambda_1, \dots, \lambda_n) + \lambda_n \lambda_{n-1} e_2(\lambda_1, \dots, \lambda_n) + \dots + \lambda_n \dots \lambda_1 e_n(\lambda_1, \dots, \lambda_n)$$

by introducing new parameters  $A_1 \geq A_2 \geq \dots \geq A_n > 0$  for non-symmetrically appearing  $\lambda_1, \dots, \lambda_n$  and commutative variables  $a_1, a_2, \dots, a_n$  for symmetrically appearing  $\lambda_1, \dots, \lambda_n$ .

$$\Psi_{a_1, a_2, \dots, a_n}^{A_1, A_2, \dots, A_n} := 1 + A_1 e_1(a_1, \dots, a_n) + A_1 A_2 e_2(a_1, \dots, a_n) + \dots + A_1 A_2 \dots A_n e_n(a_1, \dots, a_n)$$

which we abbreviate as (with  $A_{1\dots k} := A_1 A_2 \dots A_k$ )

$$\Psi_{12\dots n}^{12\dots n} = 1 + A_1 e_1 + A_{12} e_2 + \dots + A_{1\dots n} e_n$$

and proposed a conjecture (in [4], c.f. Conj. 1.5. in [9])

$$\boxed{(\Psi_{12\dots n}^{12\dots n})^{n-1} \geq \prod_{k=1}^n \Psi_{1\dots \hat{k}\dots n}^{1\dots \hat{k}\dots n}} \quad (*)$$

hold coefficientwise in the ring  $\mathbf{R}[a_1, \dots, a_n]$  of polynomials in (commuting) indeterminants  $a_1, a_2, \dots, a_n$  (and verified it with a number of refinements for  $n \leq 9$ ).

It is trivial for  $n = 2$ :

$$\Psi_{12}^{12} = a + A_1(a_1 + a_2) + A_1 A_2 a_1 a_2 \geq (1 + A_2 a_2)(1 + A_1 a_1) = \Psi_2^2 \Psi_1^1$$

because

$$\Psi_{12}^{12} - \Psi_2^2 \Psi_1^1 = (A_1 - A_2)a_2$$

(and we assumed  $A_1 \geq A_2 > 0$ ).

Note that the r.h.s. of the Conjecture is not symmetric in variables  $a_1, \dots, a_n$ , but by studying the derivatives of

$$\frac{\Psi_{1\dots n}^{1\dots n}}{\Psi_{1\dots \hat{k}\dots n}^{1\dots \hat{k}\dots n}}$$

in [4] we stated a strengthened symmetric version.

## Conjecture

Let  $A_1 \geq \dots \geq A_n \geq, a_1, \dots, a_n \geq 0$ . Then the following inequality for symmetric functions in  $a_1, \dots, a_n$

$$\Psi_{123\dots n}^{112\dots n-1} \Psi_{1234\dots n}^{1223\dots n-1} \dots \Psi_{12\dots n-2\ n-1\ n}^{12\dots n-2\ n-1\ n-1} \geq \Psi_{12\dots n-1}^{1\ 2\dots n-1} \Psi_{12\dots n-2\ n}^{1\ 2\dots n-1} \dots \Psi_{23\dots n-1}^{1\ 2\dots n-1}$$

i.e.

$$\boxed{\prod_{k=1}^{n-1} \Psi_{12\dots k\ k+1\dots n}^{1\ 2\dots k\ k\dots n} \geq \prod_{k=1}^n \Psi_{12\dots \hat{k}\dots n}^{1\ 2\dots n-1}} \quad (**)$$

holds true coefficientwise ( $m$ -positivity).

Now by the following Lemma we interpreted the Conjecture  $(**)$  as (polynomial wrt Schur functions) a Hadamard type inequality for certain non symmetric matrices.

### Lemma

For any  $k$ ,  $(1 \leq k \leq n)$ , we have

$$\Psi_{1\dots \hat{k}\dots n}^{1\dots k\dots n-1} = \sum_{j=0}^{n-1} c_j a_k^{n-1-j}$$

where

$$c_{n-1-j} = (-1)^j \sum_{i=j}^{n-1} X_1 \cdots X_i e_{i-j}, \quad j = 0, \dots, n-1.$$

By the Lemma, the right hand side of (\*\*) can be written as  $R_n = \prod_{k=1}^n \left( \sum_{j=0}^{n-1} c_j \xi_k^{n-1-j} \right)$  and can be written as

$$R_n = \begin{vmatrix} 1 & -e_1 & e_2 & -e_3 & \cdots & (-1)^n e_n \\ & 1 & -e_1 & e_2 & -e_3 & \cdots \\ & & \ddots & & & \\ c_0 & c_1 & c_2 & \cdots & c_n & \cdots \\ & c_0 & c_1 & c_2 & \cdots & c_n \\ & & \ddots & & & \\ & & & c_0 & c_1 & c_2 & \cdots & c_n \end{vmatrix} (=: \begin{vmatrix} A & B \\ C & D \end{vmatrix})$$

can be simplified as

$$= |A| \cdot |D - CA^{-1}B| = |D - CA^{-1}B|.$$

The entries of the  $n \times n$  matrix  $\Delta := D - CA^{-1}B$  are given by

$$\delta_{ij} = \begin{cases} (-1)^{j-i-1} \sum_{k=j+1}^n A_1 \cdots A_{k+i-j} e_k, & 0 \leq i < j \leq n-1 \\ (-1)^{j-i} \sum_{k=0}^j A_1 \cdots A_{k+i-j} e_k, & 0 \leq j \leq i \leq n-1 \end{cases}$$

For example, for  $n = 3$

$$\Delta_3 = \begin{vmatrix} 1 & A_1 e_2 + A_1 A_2 e_3 & -A_1 e_3 \\ -A_1 & 1 + A_1 e_1 & A_1 A_2 e_3 \\ A_1 A_2 & -A_1 - A_1 A_2 e_1 & 1 + A_1 e_1 + A_1 A_2 e_2 \end{vmatrix}$$

By elementary operations we get

$$\Delta_3 = \begin{vmatrix} 1 & * & * \\ 0 & \Psi_{123}^{112} & A_1(A_2 - A_1)e_3 \\ 0 & A_2 - A_1 & \Psi_{123}^{122} \end{vmatrix} = \begin{vmatrix} \Psi_{123}^{112} & A_1(A_2 - A_1)e_3 \\ A_2 - A_1 & \Psi_{123}^{122} \end{vmatrix}$$

Similarly, for  $n = 4$  we obtain

$$\Delta_4 = \begin{vmatrix} \Psi_{1234}^{1123} & -A_1(A_1 - A_2)e_3 - A_1A_2(A_1 - A_3)e_4 & A_1(A_1 - A_2)e_4 \\ -(A_1 - A_2) & \Psi_{1234}^{1223} & -A_1A_2(A_2 - A_3)e_4 \\ A_1(A_2 - A_3) & -(A_1 - A_3) - A_1(A_2 - A_3)e_1 & \Psi_{1234}^{1233} \end{vmatrix}$$

In general

$$\Delta_n = \det(\delta'_{ij})_{1 \leq i, j \leq n-1}$$

where

$$\delta'_{ij} = \begin{cases} (-1)^{j-i} \sum_{k=j+1}^n A_1 \cdots A_{k+i-j-1} (A_i - A_{k+i-j}) e_k, & 1 \leq i < j \leq n-1 \\ \Psi_1^1 \dots \overset{i}{\underset{j}{\dots}} \dots \overset{n}{\underset{n}{\dots}} & , i = j \\ (-1)^{j-i} \sum_{k=0}^j A_1 \cdots A_{k+i-j-1} (A_{k+i-j} - A_i) e_k & , 1 \leq j < i \leq n-1 \end{cases}$$

Then we conjecture that the following Hadamard type inequality

$$\prod_{i=1}^{n-1} \delta'_{ii} \geq \det(\delta'_{ij}), \quad (\delta'_{ij})_{1 \leq i, j \leq n-1}$$

should hold coefficientwise w.r.t. Schur functions in  $a_1, \dots, a_n$ .

Let  $a_1, \dots, a_n, A_1, \dots, A_n, n \geq 1$  be two sets of commuting indeterminates. For any  $l, 1 \leq l \leq n$  and any sequences  $1 \leq i_1 \leq \dots \leq i_l \leq n, 1 \leq j_1, \dots, j_l \leq n$  we define polynomials  $\Psi_J^I = \Psi_{j_1 \dots j_l}^{i_1 \dots i_l} \in \mathcal{Q}[a_1, \dots, a_n, A_1, \dots, A_n]$  as follows:

$$\Psi_J^I := \sum_{k=0}^l e_k(a_{j_1}, a_{j_2}, \dots, a_{j_l}) A_{i_1} A_{i_2} \cdots A_{i_k}, \quad (l \geq 1), \quad \Psi_\emptyset^\emptyset := 1 \quad (j = 0)$$

where  $e_k$  is the  $k$ -th elementary symmetric function.

The polynomials  $\Psi_J^I$  are symmetric w.r.t.  $a_{j_1}, a_{j_2}, \dots, a_{j_l}$ , but nonsymmetric w.r.t.  $A_{i_1}, A_{i_2}, \dots, A_{i_l}$ . By specializing  $A_i$ 's to assume real values such that  $A_{i_1} \geq A_{i_2} \geq \dots \geq A_{i_l} \geq 0$  then we obtain polynomials in  $a_j$ 's satisfying the following simple but important property.

### **Proposition** (Partition property)

Let  $(I_1, \dots, I_s)$  and  $(J_1, \dots, J_s)$  be ordered set partitions of respective sets  $I = \bigcup_{p=1}^s I_p$  and  $J = \bigcup_{p=1}^s J_p$  such that  $|I_p| = |J_p|, 1 \leq p \leq s$ . Then the inequality

$$\Psi_J^I \geq \prod_{p=1}^s \Psi_{J_p}^{I_p}$$

holds coefficientwise w.r.t.  $a_j$ 's.

**Proof.** Proof is evident from the definition of  $\Psi_J^I$  and the monotonicity of  $A_i$ 's. □

For the powers  $(\Psi_J^I)^m$  we made the following conjecture in ([4]):

### **Conjecture**(Weighted Multiset Partition Conjecture)

For given natural number  $m$  and sets  $I$  and  $J, |I| = |J|$ , of natural numbers let  $(I_1, \dots, I_s)$  and  $(J_1, \dots, J_s)$  be the partitions of the multiset  $I^m$  consisting of  $m$  copies of all elements of  $I$  and similarly for  $J^m$ .

(i) Then the inequality  $(\Psi_J^I)^m \geq \prod_{p=1}^s \Psi_{J_p}^{I_p}$  holds coefficientwise w.r.t.  $a_j$ 's.

(ii) The difference  $(\Psi_J^I)^m - \prod_{p=1}^s \Psi_{J_p}^{I_p}$  is multi-Schur positive with respect to partial alphabets

corresponding to the atoms of the intersection lattice of the set system  $\{J_1, \dots, J_s\}$ .

# Tropical Version of Atiyah–Sutcliffe Conjecture for almost collinear configurations ( $n = 3$ )

In case  $n = 3$  we illustrate the tropical version of Atiyah–Sutcliffe conjecture.  
Let  $n = 3$ ,  $A_1 \geq A_2 \geq A_3 > 0$ .

$$\begin{aligned} E(x) &= (1 + a_1x)(1 + a_2x)(1 + a_3x) \\ &= 1 + e_1x + e_2x^2 + e_3x^3 \end{aligned}$$

$$\begin{aligned} E^{(1)}(x) &= (1 + a_2x)(1 + a_3x) \\ &= 1 + (a_2 + a_3)x + a_2a_3x^2 = 1 + e_1^{(1)}x + e_2^{(1)}x^2 \end{aligned}$$

$$\begin{aligned} E^{(2)}(x) &= (1 + a_1x)(1 + a_3x) \\ &= 1 + (a_1 + a_3)x + a_1a_3x^2 = 1 + e_1^{(2)}x + e_2^{(2)}x^2 \end{aligned}$$

$$\begin{aligned} E^{(3)}(x) &= (1 + a_1x)(1 + a_2x) \\ &= 1 + (a_1 + a_2)x + a_1a_2x^2 = 1 + e_1^{(3)}x + e_2^{(3)}x^2 \end{aligned}$$

$$\begin{aligned}
AS_3 &= \Psi_{123}^{123}\Psi_{123}^{123} - \Psi_{12}^{12}\Psi_{13}^{13}\Psi_{23}^{23} \\
&= (1 + A_1 e_1 + A_1 A_2 e_2 + A_1 A_2 A_3 e_3)^2 - \left(1 + A_2 e_1^{(1)} + A_2 A_3 e_2^{(1)}\right) \cdot \\
&\quad \cdot \left(1 + A_1 e_1^{(2)} + A_1 A_3 e_2^{(2)}\right) \cdot \\
&\quad \cdot \left(1 + A_1 e_1^{(3)} + A_1 A_2 e_2^{(3)}\right)
\end{aligned}$$

$$\begin{aligned}
AS_3^{tr op} &= \begin{bmatrix} t^1 \end{bmatrix} \text{subs} (A_1 = e^{\alpha_1 t}, A_2 = e^{\alpha_2 t}, A_3 = e^{\alpha_3 t}, AS_3) \\
&= \begin{bmatrix} t^1 \end{bmatrix} \text{subs} (A_1 = 1 + \alpha_1 t, A_2 = 1 + \alpha_2 t, A_3 = 1 + \alpha_3 t, AS_3) \\
&= \begin{bmatrix} t^1 \end{bmatrix} \left[ (1 + (1 + \alpha_1 t)e_1 + (1 + (\alpha_1 + \alpha_2)t)e_2 + (1 + (\alpha_1 + \alpha_2 + \alpha_3)t)e_3)^2 - \right. \\
&\quad - \left(1 + (1 + \alpha_2 t)e_1^{(1)} + (1 + (\alpha_2 + \alpha_3)t)e_2^{(1)}\right) \cdot \\
&\quad \cdot \left(1 + (1 + \alpha_1 t)e_1^{(2)} + (1 + (\alpha_1 + \alpha_3)t)e_2^{(2)}\right) \cdot \\
&\quad \cdot \left(1 + (1 + \alpha_1 t)e_1^{(3)} + (1 + (\alpha_1 + \alpha_2)t)e_2^{(3)}\right) \left] \right. \\
&= \begin{bmatrix} t^1 \end{bmatrix} \left[ (1 + e_1 + e_2 + e_3 + (\alpha_1 e_1 + (\alpha_1 + \alpha_2)e_2 + (\alpha_1 + \alpha_2 + \alpha_3)e_3)t)^2 \right. \\
&\quad - \left(1 + e_1^{(1)} + e_2^{(1)} + (\alpha_2 e_1^{(1)} + (\alpha_2 + \alpha_3)e_2^{(1)})t\right) \cdot \\
&\quad \cdot \left(1 + e_1^{(2)} + e_2^{(2)} + (\alpha_1 e_1^{(2)} + (\alpha_1 + \alpha_3)e_2^{(2)})t\right) \cdot \\
&\quad \cdot \left(1 + e_1^{(3)} + e_2^{(3)} + (\alpha_1 e_1^{(3)} + (\alpha_1 + \alpha_2)e_2^{(3)})t\right) \left] = \right.
\end{aligned}$$

$$\begin{aligned}
&= \left[ t^1 \right] \left[ \left( E(1) + (\alpha_1 e_1 + (\alpha_1 + \alpha_2)e_2 + (\alpha_1 + \alpha_2 + \alpha_3)e_3 t)^2 \right) \right. \\
&\quad - \left( E^{(1)}(1) + \left( \alpha_2 e_1^{(1)} + (\alpha_2 + \alpha_3)e_2^{(1)} \right) t \right) \cdot \\
&\quad \cdot \left( E^{(2)}(1) + \left( \alpha_1 e_1^{(2)} + (\alpha_1 + \alpha_3)e_2^{(2)} \right) t \right) \cdot \\
&\quad \cdot \left. \left( E^{(3)}(1) + \left( \alpha_1 e_1^{(3)} + (\alpha_1 + \alpha_2)e_2^{(3)} \right) t \right) \right] = \\
&= 2E(1)(\alpha_1 e_1 + (\alpha_1 + \alpha_2)e_2 + (\alpha_1 + \alpha_2 + \alpha_3)e_3 t)^2 \\
&\quad - E^{(2)}(1)E^{(2)}(1) \left( \alpha_2 e_1^{(1)} + (\alpha_2 + \alpha_3)e_2^{(1)} \right) \\
&\quad - E^{(1)}(1)E^{(3)}(1) \left( \alpha_1 e_1^{(2)} + (\alpha_1 + \alpha_3)e_2^{(2)} \right) \\
&\quad - E^{(1)}(1)E^{(2)}(1) \left( \alpha_1 e_1^{(3)} + (\alpha_1 + \alpha_2)e_2^{(3)} \right) =
\end{aligned}$$

Now we use  $E^{(2)}(1)E^{(3)}(1) = (1 + a_1)(1 + a_3)(1 + a_1)(1 + a_2) = E(1)(1 + a_1)$  etc.

$$\begin{aligned}
&= E(1) \left[ 2\alpha_1 e_1 + (2\alpha_1 + 2\alpha_2)e_2 + (2\alpha_1 + 2\alpha_2 + 2\alpha_3)e_3 \right. \\
&\quad - (1 + a_1) \left( \alpha_2 e_1^{(1)} + (\alpha_2 + \alpha_3)e_2^{(1)} \right) \\
&\quad - (1 + a_2) \left( \alpha_1 e_1^{(2)} + (\alpha_1 + \alpha_3)e_2^{(2)} \right) \\
&\quad \left. - (1 + a_3) \left( \alpha_1 e_1^{(3)} + (\alpha_1 + \alpha_2)e_2^{(3)} \right) \right] =
\end{aligned}$$

By using basic relation between elementary symmetric polynomials

$$e_k(a_1, \dots, a_n) = a_j e_{k-1}(a_1, \dots, \widehat{a_j}, \dots, a_n) + e_k(a_1, \dots, \widehat{a_j}, \dots, a_n)$$

$$\begin{aligned} &= E(1) \left\{ 2\alpha_1 e_1 + (2\alpha_1 + 2\alpha_2)e_2 + (2\alpha_1 + 2\alpha_2 + 2\alpha_3)e_3 \right. \\ &\quad - \left[ \alpha_2 e_1^{(1)} + \alpha_2 e_2 - \cancel{\alpha_2 e_2^{(1)}} + (\underline{\alpha_2} + \alpha_3) \underline{e_2^{(1)}} + (\alpha_2 + \alpha_3) e_3 \right] \\ &\quad - \left[ \alpha_1 e_1^{(2)} + \alpha_1 e_2 - \cancel{\alpha_1 e_2^{(2)}} + (\underline{\alpha_1} + \alpha_3) \underline{e_2^{(2)}} + (\alpha_1 + \alpha_3) e_3 \right] \\ &\quad \left. - \left[ \alpha_1 e_1^{(3)} + \alpha_1 e_2 - \cancel{\alpha_1 e_2^{(3)}} + (\underline{\alpha_1} + \alpha_2) \underline{e_2^{(3)}} + (\alpha_1 + \alpha_2) e_3 \right] \right\} = \end{aligned}$$

By using  $2e_1 = e_1^{(1)} + e_1^{(2)} + e_1^{(3)}$ ,  $e_2 = e_2^{(1)} + e_2^{(2)} + e_2^{(3)}$

$$= E(1) \left[ (\alpha_1 - \alpha_2) e_1^{(1)} + \alpha_2 e_2 - (\alpha_3 e_2^{(1)} + \alpha_3 e_2^{(2)} + \alpha_2 e_2^{(3)}) \right]$$

$$AS_3^{trop} = E(1) \left[ (\alpha_1 - \alpha_2) e_1^{(1)} + (\alpha_2 - \alpha_3) (e_2^{(1)} + e_2^{(2)}) \right]$$

$$(E(1) = 1 + e_1 + e_2 + e_3)$$

In general it would be

$$AS_n^{trop} = (E(1))^{n-2} \left[ (\alpha_1 - \alpha_2)e_1^{(1)} + \cdots + (\alpha_{n-1} - \alpha_n) \left( e_{n-1}^{(1)} + \cdots + e_{n-1}^{(n-1)} \right) \right]$$

# Example Verification supporting our Multiset Partition Conjecture Using Maple (1/4)

Let  $a_1, a_2, \dots, a_6$  be commuting variables and let  $A_1 \geq A_2 \geq \dots \geq A_6 > 0$  be nonnegative real numbers.

Let  $f_1 = a_2 + a_4 + a_5$ ,  $f_2 = a_2 \cdot a_4 + a_2 \cdot a_5 + a_4 \cdot a_5$ ,  $f_3 = a_2 \cdot a_4 \cdot a_5$ ,  $h_1 = a_2 + a_5$ ,  $h_2 = a_2 \cdot a_5$ , and  $e_1 = a_1 + a_3 + a_6$ ,  $e_2 = a_1 \cdot a_3 + a_1 \cdot a_6 + a_3 \cdot a_6$ ,  $e_3 = a_1 \cdot a_3 \cdot a_6$  be the elementary symmetric functions of the alphabets  $a_2, a_4, a_5$  and  $a_2, a_5$  and  $a_1, a_3, a_6$ .

The  $\psi_{123456}$  function of the original alphabet  $a_1, a_2, a_3, a_4, a_5, a_6$ , is

$$\begin{aligned} \psi_{123456} = & \\ 1 + A[1](e_1 + f_1) + A[1]A[2] \cdot (e_2 + e_1 \cdot f_1 + f_2) + A[1]A[2]A[3](e_3 + e_2 \cdot f_1 + e_1 \cdot f_2 + f_3) + A[1]A[2]A[3]A[4] & \\ + A[1]A[2]A[3]A[4]A[5]A[6]e_3 \cdot f_3 (= \psi_{123456}) & \end{aligned}$$

The  $\psi_{1245}$  function of the alphabet  $a_2, a_4, a_5$  is

$$\psi_{1245} = 1 + A[2] \cdot f_1 + A[2] \cdot A[4] \cdot f_2 + A[2] \cdot A[4] \cdot A[5] \cdot f_3 (= D_1).$$

The  $\psi_{136}$  function of the alphabet  $a_1, a_3, a_6$  is

$$\psi_{136} = 1 + A[1] \cdot e_1 + A[1] \cdot A[2] \cdot e_2 + A[1] \cdot A[3] \cdot A[6] \cdot e_3.$$

The  $\psi_{1346}$  function of the alphabet  $a_1, a_3, a_4, a_6$  is

$$\psi_{1346} = 1 + A[1](e_1 + a[4]) + A[1]A[3](e_2 + e_1 a[4]) + A[1]A[3]A[4](e_3 + e_2 a[4]) + A[1]A[3]A[4]A[6]e_3 a[4] (= d_1).$$

The  $\psi_{12356}$  function of the alphabet  $a_1, a_2, a_3, a_5, a_6$  is

$$\begin{aligned} \psi_{12356} = & \\ (1 + A[1] \cdot (e_1 + h_1) + A[1] \cdot A[2] \cdot (e_2 + e_1 \cdot h_1 + h_2) + A[1] \cdot A[2] \cdot A[3] \cdot (e_3 + e_2 \cdot h_1 + e_1 \cdot h_2) + & \\ + A[1] \cdot A[2] \cdot A[3] \cdot A[5] \cdot (e_3 \cdot h_1 + e_2 \cdot h_2) + A[1] \cdot A[2] \cdot A[3] \cdot A[5] \cdot A[6] \cdot e_3 \cdot h_2) (= d_2). & \end{aligned}$$

Then our Weighted Multiset Partition Conjecture (Conjecture 3.2 in arxiv.org.math.0609174.pdf) reads as the following coefficientwise inequality:

$$\psi_{123456}^2 \geq \psi_{1245} \cdot \psi_{1346} \cdot \psi_{12356}.$$



## Example Verification supporting our Multiset Partition Conjecture Using Maple (2/4)

The final formula for

$$(\psi_{123456})^2 - \psi_{1245} \cdot \psi_{1346} \cdot \psi_{12356}$$

in terms of Schur functions  $t_1, t_{11}, t_{111}, t_2, t_{21}, t_{211}, t_{22}, t_{221}, t_{222}$  of  $a[1], a[3], a[6]$  and Schur functions  $s[1], s[2], s[1, 1], s[2, 1], s[2, 2], s[1, 1, 1], s[2, 1, 1], s[2, 2, 1], s[2, 2, 2]$  of  $a[2], a[4], a[5]$  is  $Z6c$  below (and its coefficients are positive).

```
Z6c :=  
sort(map(factor, collect(Z6b, {t1, t11, t111, t2, t21, t211, t22, t221, t222, s[1], s[2], s[1, 1], s[2, 1], s[2, 2],  
s[1, 1, 1], s[2, 1, 1], s[2, 2, 1], s[2, 2, 2]}), distributed)), {A[1], A[2], A[3], A[4], A[5], A[6]});
```

# Example Verification supporting our Multiset Partition Conjecture Using Maple (3/4)

$$\begin{aligned} Z6c := & (A[2] - A[3]) \cdot s[1, 1] \cdot t22 \cdot A[1]^2 \cdot A[2] \cdot A[3] \cdot A[4] + (A[2] \cdot A[3] - A[2] \cdot A[4] + A[2] \cdot A[5] \\ & - A[4] \cdot A[6]) \cdot s[2, 2] \cdot t111 \cdot A[1]^2 \cdot A[2] \cdot A[3] \cdot A[4] + (A[2] - A[3]) \cdot t22 \cdot A[1]^2 \cdot A[2] + (2 \cdot A[2] \cdot A[3] \\ & - A[2] \cdot A[4] - A[4] \cdot A[5]) \cdot s[2, 2, 1] \cdot A[1]^2 \cdot A[2] + (A[2] - A[4]) \cdot s[2, 2] \cdot A[1]^2 \cdot A[2] \\ & + (A[1] - A[4]) \cdot s[2, 1] \cdot A[1] \cdot A[2] + (2 \cdot A[1] \cdot A[3] - A[1] \cdot A[4] - A[4] \cdot A[5]) \cdot s[2, 1, 1] \cdot A[1] \cdot A[2] \\ & + (2 \cdot A[1] \cdot A[2] - A[1] \cdot A[3] - A[2]^2 + 2 \cdot A[2] \cdot A[3] - 2 \cdot A[2] \cdot A[4]) \cdot s[1, 1] \cdot t1 \cdot A[1] \\ & + (A[1] - A[2]) \cdot s[2] \cdot A[1] + (2 \cdot A[1] \cdot A[2]^2 + A[1] \cdot A[2] \cdot A[3] - A[1] \cdot A[2] \cdot A[4] - A[1] \cdot A[3] \cdot A[4] \\ & - A[2]^2 \cdot A[3] - A[2]^2 \cdot A[4] + A[2] \cdot A[3] \cdot A[4]) \cdot s[1, 1] \cdot t11 \cdot A[1] + (A[1] - A[2]) \cdot s[1] \cdot t1 \cdot A[1] \\ & + (2 \cdot A[1] \cdot A[2] - A[1] \cdot A[3] - A[2]^2) \cdot s[1] \cdot t11 \cdot A[1] + (4 \cdot A[1] \cdot A[2]^2 \cdot A[3] - A[1] \cdot A[2]^2 \cdot A[4] \\ & - A[1] \cdot A[2] \cdot A[3]^2 + A[1] \cdot A[2] \cdot A[3] \cdot A[4] - A[1] \cdot A[2] \cdot A[3] \cdot A[5] - A[1] \cdot A[3] \cdot A[4] \cdot A[6] \\ & - A[2]^2 \cdot A[3] \cdot A[4] - A[2]^2 \cdot A[3] \cdot A[5] - A[2] \cdot A[3] \cdot A[4]^2 + 2 \cdot A[2] \cdot A[3] \cdot A[4] \cdot A[5]) \cdot s[1, 1] \\ & \cdot t111 \cdot A[1] + (2 \cdot A[1] \cdot A[3] - A[1] \cdot A[4] - A[4] \cdot A[5]) \cdot s[1, 1, 1] \cdot A[2] + (A[1] \cdot A[2]^2 \\ & + A[1] \cdot A[2] \cdot A[3] - A[1] \cdot A[3] \cdot A[4] - A[2]^2 \cdot A[3] + A[2] \cdot A[3] \cdot A[4] - A[2] \cdot A[3] \cdot A[5]) \cdot s[1] \cdot t111 \\ & \cdot A[1] + (A[5] - A[6]) \cdot s[2, 2] \cdot t222 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \cdot A[4]^2 \cdot A[5] + (A[5] - A[6]) \cdot s[2, 2, 1] \\ & \cdot t221 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \cdot A[4]^2 \cdot A[5] + (A[5] - A[6]) \cdot s[2, 2, 1] \cdot t222 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \\ & \cdot A[4]^2 \cdot A[5] \cdot A[6] + (A[4] - A[5]) \cdot s[2, 2, 2] \cdot t211 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \cdot A[4] \cdot A[5] \cdot A[6] \\ & \dots \\ & + (A[2] - A[5]) \cdot s[1] \cdot t222 \cdot A[1]^2 \cdot A[2] \cdot A[3]^2 \cdot A[4] + (2 \cdot (A[3] - A[5])) \cdot s[2, 2, 2] \cdot t1 \cdot A[1]^2 \\ & \cdot A[2]^2 \cdot A[3] \cdot A[4] + (A[3] \cdot A[5] - A[4] \cdot A[6]) \cdot s[2, 2] \cdot t211 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3] \cdot A[4] \\ & + (2 \cdot A[3] \cdot A[4] \cdot A[5] - A[3] \cdot A[4] \cdot A[6] - A[3] \cdot A[5]^2 + A[3] \cdot A[5] \cdot A[6] - A[4] \cdot A[5] \cdot A[6]) \cdot s[2, 2, 1] \\ & \cdot t211 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3] \cdot A[4] + (A[3] - A[5]) \cdot s[2, 2, 1] \cdot t2 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3] \cdot A[4] \\ & + (A[3] - A[4]) \cdot s[2, 2] \cdot t21 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3] \cdot A[4] + (A[3] - A[5]) \cdot s[2, 2, 1] \cdot t21 \cdot A[1]^2 \cdot A[2]^2 \\ & \cdot A[3] \cdot A[4]^2 + (2 \cdot A[4] \cdot A[5] - A[4] \cdot A[6] - A[5]^2) \cdot s[2, 1, 1] \cdot t221 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \cdot A[4] \\ & + (A[4] - A[6]) \cdot s[2, 1] \cdot t221 \cdot A[1]^2 \cdot A[2]^2 \cdot A[3]^2 \cdot A[4] \end{aligned}$$

> nops(Z6c);

86

# Example Verification supporting our Multiset Partition Conjecture Using Maple (4/4)

```
for k to 86 do k, op(k, Z6c) end do
```

```
1, ( $A[5] - A[6]$ ) $s[2, 2, 1]t222A[1]^2 A[2]^2 A[3]^2 A[4]^2 A[5]A[6]$ 
2, ( $A[5] - A[6]$ ) $s[2, 2, 1]t221A[1]^2 A[2]^2 A[3]^2 A[4]^2 A[5]$ 
3, ( $A[5] - A[6]$ ) $s[2, 2]t222A[1]^2 A[2]^2 A[3]^2 A[4]^2 A[5]$ 
4, ( $A[4] - A[5]$ ) $s[2, 2, 2]t211A[1]^2 A[2]^2 A[3]^2 A[4]A[5]A[6]$ 
5, ( $A[4] - A[6]$ ) $s[2, 1, 1]t222A[1]^2 A[2]^2 A[3]^2 A[4]A[5]A[6]$ 
6, ( $A[5] - A[6]$ ) $s[2, 2]t221A[1]^2 A[2]^2 A[3]^2 A[4]^2$ 
7, ( $A[4] - A[6]$ ) $s[2, 1]t222A[1]^2 A[2]^2 A[3]^2 A[4]A[5]$ 
8, ( $A[4] - A[5]$ ) $s[2, 2, 2]t21A[1]^2 A[2]^2 A[3]^2 A[4]A[5]$ 
9, ( $A[2] - A[6]$ ) $s[1, 1, 1]t222A[1]^2 A[2]A[3]^2 A[4]A[5]A[6]$ 
10, ( $A[4] - A[5]$ ) $s[2, 2, 2]t2A[1]^2 A[2]^2 A[3]^2 A[4]$ 
11, ( $A[4] - A[5]$ ) $s[2]t222A[1]^2 A[2]^2 A[3]^2 A[4]$ 
12, ( $A[4] - A[6]$ ) $s[2, 1]t221A[1]^2 A[2]^2 A[3]^2 A[4]$ 
13, ( $A[4]A[5] - A[4]A[6] - A[5]^2$ ) $s[2, 1, 1]t221A[1]^2 A[2]^2 A[3]^2 A[4]$ 
...
83, ( $A[1]A[2]^2 + A[1]A[2]A[3] - A[1]A[3]A[4] - A[2]^2 A[3] + A[2]A[3]A[4] - A[2]A[3]A[5]$ ) $s[1]t111A[1]$ 
84, ( $2A[1]A[3] - A[1]A[4] - A[4]A[5]$ ) $s[1, 1, 1]A[2]$ 
85, ( $A[1] - A[4]$ ) $s[1, 1]A[2]$ 
86, ( $A[1] - A[2]$ ) $s[1]$ 
```

# Tropical Version of Atiyah–Sutcliffe Conjecture for almost collinear configurations ( $n = 4$ )

Tropical version for  $n = 4$  is reduced analogously to the following expression

$$\begin{aligned} AS_4^{trop} &= E(1)^2 [3\alpha_1 e_1 + 3(\alpha_1 + \alpha_2)e_2 + 3(\alpha_1 + \alpha_2 + \alpha_3)e_3 + 3(\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)e_4 \\ &\quad - (\alpha_2 e_1^{(1)} + \alpha_2 e_2 + \alpha_3 e_2^{(1)} + (\alpha_2 + \alpha_3)e_3 + \alpha_4 e_3^{(1)} + (\alpha_2 + \alpha_3 + \alpha_4)e_4) \\ &\quad - (\alpha_1 e_1^{(2)} + \alpha_1 e_2 + \alpha_3 e_2^{(2)} + (\alpha_1 + \alpha_3)e_3 + \alpha_4 e_3^{(2)} + (\alpha_1 + \alpha_3 + \alpha_4)e_4) \\ &\quad - (\alpha_1 e_1^{(3)} + \alpha_1 e_2 + \alpha_2 e_2^{(3)} + (\alpha_1 + \alpha_2)e_3 + \alpha_4 e_3^{(3)} + (\alpha_1 + \alpha_2 + \alpha_4)e_4) \\ &\quad - (\alpha_1 e_1^{(4)} + \alpha_1 e_2 + \alpha_2 e_2^{(4)} + (\alpha_1 + \alpha_2)e_3 + \alpha_3 e_3^{(4)} + (\alpha_1 + \alpha_2 + \alpha_3)e_4)] \\ &= E(1)^2 [3\alpha_1 e_1 + 3(\alpha_1 + \alpha_2)e_2 + 3(\alpha_1 + \alpha_2 + \alpha_3)e_3 + 3(\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)e_4 \\ &\quad - \underbrace{\left( \alpha_1 (e_1^{(1)} + e_1^{(2)} + e_1^{(3)} + e_1^{(4)}) + (\alpha_2 - \alpha_1)e_1^{(1)} + 3(\alpha_1 + \alpha_2)e_2 + (\alpha_3 - \alpha_2)(e_2^{(1)} + e_2^{(2)}) + \right.} \\ &\quad \left. \left. + (3(\alpha_1 + \alpha_2) + 2\alpha_3 + \alpha_3)e_3 + (\alpha_4 - \alpha_3)(e_3^{(1)} + e_3^{(2)} + e_3^{(3)}) \right) \right] \end{aligned}$$

$$AS_4^{trop} = E(1)^2 [(\alpha_1 - \alpha_2)e_1^{(1)} + (\alpha_2 - \alpha_3)(e_2^{(1)} + e_2^{(2)}) + (\alpha_3 - \alpha_4)(e_3^{(1)} + e_3^{(2)} + e_3^{(3)})]$$

Here we used the elementary formulas:

$$e_1^{(1)} + e_1^{(2)} + e_1^{(3)} + e_1^{(4)} = 3e_1, \quad e_2^{(1)} + e_2^{(2)} + e_2^{(3)} + e_2^{(4)} = 2e_2, \quad e_3^{(1)} + e_3^{(2)} + e_3^{(3)} + e_3^{(4)} = e_3.$$

# References

- [1] Atiyah M., Sutcliffe P., The Geometry of Point Particles. arXiv: hep-th/0105179 (32 pages). Proc.R.Soc.Lond. A (2002) 458, 1089-115.
- [2] Atiyah M., Sutcliffe P., Polyhedra in Physics, Chemistry and Geometry, arXiv: math-ph/03030701 (22 pages), "Milan J.Math." 71:33-58 (2003)
- [3] Eastwood M., Norbury P., A proof of Atiyah's conjecture on configurations of four points in Euclidean three space, Geometry and Topology 5(2001) 885-893.
- [4] Svrtan D., Urbija I., Verification and Strengthening of the Atiyah-Sutcliffe Conjectures for Several Types of Configurations, arXiv: math/0609174 (49 pages).
- [5] Atiyah M., An Unsolved Problem in Elementary Geometry , [www.math.missouri.edu/archive/Miller-Lectures/atiyah/atiyah.html](http://www.math.missouri.edu/archive/Miller-Lectures/atiyah/atiyah.html).
- [6] Mazur M., Petrenko B.V., On the conjectures of Atiyah and Sutcliffe , arXiv:math.AG/1102.46612
- [7] Svrtan D., [http://bib.irb.hr/datoteka/553817.SVRTAN\\_MCC2011\\_HYPERBOLIC\\_EASTWOOD\\_NORBURY\\_FORMULA\\_FOR\\_ATIYAH\\_DETERMINANT.ppt](http://bib.irb.hr/datoteka/553817.SVRTAN_MCC2011_HYPERBOLIC_EASTWOOD_NORBURY_FORMULA_FOR_ATIYAH_DETERMINANT.ppt)
- [8] Svrtan D., Urbija I., [http://icm2006.org/v\\_f/AbsDef/Global/Posters05.pdf](http://icm2006.org/v_f/AbsDef/Global/Posters05.pdf)
- [9] Svrtan D., Urbija I., Verification and Strengthening of the Atiyah–Sutcliffe Conjectures for Several Types of Almost Collinear Configurations in Euclidean and Hyperbolic Plane, Proceedings of the 2nd Croatian Combinatorial Days. Editors: Tomislav Došlić, Ivica Martinjak, Publisher: Faculty of Civil Engineering, University of Zagreb, ISBN: 978-953-8168-33-8, Published online: July 2019., pp.143-181
- [10] Svrtan D., A proof of all three Euclidean Atiyah Sutcliffe 4-point conjectures, International Congress of Mathematicians, Seoul 2014, Abstracts, Short Communications, Poster Sessions, 136–137.
- [11] Svrtan D., <http://www.emis.de/journals/SLC/wpapers/s73vortrag/svrtan.pdf>
- [12] Malkoun J., Configurations of Points and the Symplectic Berry-Robbins Problem, Symmetry Integrability and Geometry Methods and Applications, SIGMA 10 (2014), 112, 6 pages
- [13] Malkoun J., Determinants, Choices and Combinatorics, Discrete Mathematics 342 (1), 250-255, 2019.

THANK YOU

> **INTRINSIC FORMULA FOR THE HYPERBOLIC 4  
- POINT ATIYAH DETERMINANT** 20220921

> with(LinearAlgebra) :

> **for** j **to** 4 **do** p[j] :=  $\frac{\text{product}(z - t[j, k], k = 1 .. 4)}{z - t[j, j]}$  **od**  

$$\begin{aligned} p_1 &:= (z - t_{1,2})(z - t_{1,3})(z - t_{1,4}) \\ p_2 &:= (z - t_{2,1})(z - t_{2,3})(z - t_{2,4}) \\ p_3 &:= (z - t_{3,1})(z - t_{3,2})(z - t_{3,4}) \\ p_4 &:= (z - t_{4,1})(z - t_{4,2})(z - t_{4,3}) \end{aligned} \quad (1)$$

> M[4] := map(expand, Transpose(Matrix(4, 4, [seq([seq(coeff(p[j], z, 4 - k), j = 1 .. 4)], k = 1 .. 4)])))

$$M_4 := \begin{bmatrix} 1 & -t_{1,2} - t_{1,3} - t_{1,4} & t_{1,2}t_{1,3} + t_{1,4}t_{1,2} + t_{1,4}t_{1,3} & -t_{1,2}t_{1,3}t_{1,4} \\ 1 & -t_{2,1} - t_{2,3} - t_{2,4} & t_{2,1}t_{2,3} + t_{2,4}t_{2,1} + t_{2,4}t_{2,3} & -t_{2,1}t_{2,3}t_{2,4} \\ 1 & -t_{3,1} - t_{3,2} - t_{3,4} & t_{3,1}t_{3,2} + t_{3,4}t_{3,1} + t_{3,4}t_{3,2} & -t_{3,1}t_{3,2}t_{3,4} \\ 1 & -t_{4,1} - t_{4,2} - t_{4,3} & t_{4,1}t_{4,2} + t_{4,3}t_{4,1} + t_{4,3}t_{4,2} & -t_{4,1}t_{4,2}t_{4,3} \end{bmatrix} \quad (2)$$

> Δ4 := Determinant(M[4]) : prod := product(product(t[i, k] - t[k, i], i = 1 .. k - 1), k = 1 .. 4)

$$\begin{aligned} prod &:= (t_{1,2} - t_{2,1})(t_{1,3} - t_{3,1})(t_{2,3} - t_{3,2})(t_{1,4} - t_{4,1})(t_{2,4} - t_{4,2})(t_{3,4} \\ &\quad - t_{4,3}) \end{aligned} \quad (3)$$

>

$$> D4 := \frac{\Delta 4}{prod};$$

$$\begin{aligned} D4 &:= (t_{2,1}t_{4,1}t_{4,2}t_{4,3}t_{3,1}t_{3,2} + t_{2,1}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,1} + t_{2,1}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,2} \\ &\quad - t_{2,1}t_{3,1}t_{3,2}t_{3,4}t_{4,1}t_{4,2} - t_{2,1}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,1} - t_{2,1}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,2} \\ &\quad + t_{2,3}t_{4,1}t_{4,2}t_{4,3}t_{3,1}t_{3,2} + t_{2,3}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,1} + t_{2,3}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,2} \\ &\quad - t_{2,3}t_{3,1}t_{3,2}t_{3,4}t_{4,1}t_{4,2} - t_{2,3}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,1} - t_{2,3}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,2} \\ &\quad + t_{2,4}t_{4,1}t_{4,2}t_{4,3}t_{3,1}t_{3,2} + t_{2,4}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,1} + t_{2,4}t_{4,1}t_{4,2}t_{4,3}t_{3,4}t_{3,2} \\ &\quad - t_{2,4}t_{3,1}t_{3,2}t_{3,4}t_{4,1}t_{4,2} - t_{2,4}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,1} - t_{2,4}t_{3,1}t_{3,2}t_{3,4}t_{4,3}t_{4,2} \\ &\quad + t_{3,1}t_{2,1}t_{2,3}t_{2,4}t_{4,1}t_{4,2} + t_{3,1}t_{2,1}t_{2,3}t_{2,4}t_{4,3}t_{4,1} + t_{3,1}t_{2,1}t_{2,3}t_{2,4}t_{4,3}t_{4,2} \\ &\quad - t_{3,1}t_{4,1}t_{4,2}t_{4,3}t_{2,1}t_{2,3} - t_{3,1}t_{4,1}t_{4,2}t_{4,3}t_{2,4}t_{2,1} - t_{3,1}t_{4,1}t_{4,2}t_{4,3}t_{2,4}t_{2,3} \\ &\quad + t_{3,2}t_{2,1}t_{2,3}t_{2,4}t_{4,1}t_{4,2} + t_{3,2}t_{2,1}t_{2,3}t_{2,4}t_{4,3}t_{4,1} + t_{3,2}t_{2,1}t_{2,3}t_{2,4}t_{4,3}t_{4,2} \\ &\quad - t_{3,2}t_{4,1}t_{4,2}t_{4,3}t_{2,1}t_{2,3} - t_{3,2}t_{4,1}t_{4,2}t_{4,3}t_{2,4}t_{2,1} - t_{3,2}t_{4,1}t_{4,2}t_{4,3}t_{2,4}t_{2,3} \end{aligned} \quad (4)$$



$$\begin{aligned}
& + t_{4,2} t_{2,1} t_{2,3} t_{2,4} t_{1,2} t_{1,3} + t_{4,2} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,2} + t_{4,2} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,3} \\
& - t_{4,2} t_{1,2} t_{1,3} t_{1,4} t_{2,1} t_{2,3} - t_{4,2} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,1} - t_{4,2} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,3} \\
& + t_{4,3} t_{2,1} t_{2,3} t_{2,4} t_{1,2} t_{1,3} + t_{4,3} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,2} + t_{4,3} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,3} \\
& - t_{4,3} t_{1,2} t_{1,3} t_{1,4} t_{2,1} t_{2,3} - t_{4,3} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,1} - t_{4,3} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,3} \\
& - t_{1,2} t_{3,1} t_{3,2} t_{3,4} t_{2,1} t_{2,3} - t_{1,2} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,1} - t_{1,2} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,3} \\
& + t_{1,2} t_{2,1} t_{2,3} t_{2,4} t_{3,1} t_{3,2} + t_{1,2} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,1} + t_{1,2} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,2} \\
& - t_{1,3} t_{3,1} t_{3,2} t_{3,4} t_{2,1} t_{2,3} - t_{1,3} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,1} - t_{1,3} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,3} \\
& + t_{1,3} t_{2,1} t_{2,3} t_{2,4} t_{3,1} t_{3,2} + t_{1,3} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,1} + t_{1,3} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,2} \\
& - t_{1,4} t_{3,1} t_{3,2} t_{3,4} t_{2,1} t_{2,3} - t_{1,4} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,1} - t_{1,4} t_{3,1} t_{3,2} t_{3,4} t_{2,4} t_{2,3} \\
& + t_{1,4} t_{2,1} t_{2,3} t_{2,4} t_{3,1} t_{3,2} + t_{1,4} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,1} + t_{1,4} t_{2,1} t_{2,3} t_{2,4} t_{3,4} t_{3,2} \\
& - t_{2,1} t_{1,2} t_{1,3} t_{1,4} t_{3,1} t_{3,2} - t_{2,1} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,1} - t_{2,1} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,2} \\
& + t_{2,1} t_{3,1} t_{3,2} t_{3,4} t_{1,2} t_{1,3} + t_{2,1} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,2} + t_{2,1} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,3} \\
& - t_{2,3} t_{1,2} t_{1,3} t_{1,4} t_{3,1} t_{3,2} - t_{2,3} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,1} - t_{2,3} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,2} \\
& + t_{2,3} t_{3,1} t_{3,2} t_{3,4} t_{1,2} t_{1,3} + t_{2,3} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,2} + t_{2,3} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,3} \\
& - t_{2,4} t_{1,2} t_{1,3} t_{1,4} t_{3,1} t_{3,2} - t_{2,4} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,1} - t_{2,4} t_{1,2} t_{1,3} t_{1,4} t_{3,4} t_{3,2} \\
& + t_{2,4} t_{3,1} t_{3,2} t_{3,4} t_{1,2} t_{1,3} + t_{2,4} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,2} + t_{2,4} t_{3,1} t_{3,2} t_{3,4} t_{1,4} t_{1,3} \\
& - t_{3,1} t_{2,1} t_{2,3} t_{2,4} t_{1,2} t_{1,3} - t_{3,1} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,2} - t_{3,1} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,3} \\
& + t_{3,1} t_{1,2} t_{1,3} t_{1,4} t_{2,1} t_{2,3} + t_{3,1} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,1} + t_{3,1} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,3} \\
& - t_{3,2} t_{2,1} t_{2,3} t_{2,4} t_{1,2} t_{1,3} - t_{3,2} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,2} - t_{3,2} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,3} \\
& + t_{3,2} t_{1,2} t_{1,3} t_{1,4} t_{2,1} t_{2,3} + t_{3,2} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,1} + t_{3,2} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,3} \\
& - t_{3,4} t_{2,1} t_{2,3} t_{2,4} t_{1,2} t_{1,3} - t_{3,4} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,2} - t_{3,4} t_{2,1} t_{2,3} t_{2,4} t_{1,4} t_{1,3} \\
& + t_{3,4} t_{1,2} t_{1,3} t_{1,4} t_{2,1} t_{2,3} + t_{3,4} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,1} + t_{3,4} t_{1,2} t_{1,3} t_{1,4} t_{2,4} t_{2,3} \Big) / ((t_{1,2} \\
& - t_{2,1}) (t_{1,3} - t_{3,1}) (t_{2,3} - t_{3,2}) (t_{1,4} - t_{4,1}) (t_{2,4} - t_{4,2}) (t_{3,4} - t_{4,3}))
\end{aligned}$$

> nops(num(D4)) (5)

216

> denom(D4) (6)

$$(t_{2,3} - t_{3,2}) (t_{2,4} - t_{4,2}) (t_{1,3} - t_{3,1}) (t_{1,4} - t_{4,1}) (t_{1,2} - t_{2,1}) (t_{3,4} - t_{4,3})$$

>

> D4a := map(factor, collect(factor(expand(subs(seq(seq(t[i,j] =  $\frac{(s \cdot A[i,j] + B[i,j])}{2}$ , i = 1 .. j - 1), j = 1 .. 4), seq(seq(t[j,i] =  $\frac{(s \cdot A[i,j] - B[i,j])}{2}$ , i = 1 .. j - 1), j = 1 .. 4)), D4)), s));

> indets(D4a) (7)

$$\{s, A_{1,2}, A_{1,3}, A_{1,4}, A_{2,3}, A_{2,4}, A_{3,4}, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}\}$$

> expand(cot(A + B)) (8)

$$\frac{-1 + \cot(A) \cot(B)}{\cot(A) + \cot(B)}$$

>  $\cos\left(\frac{\alpha}{2}\right)^2 :$

>

>  $cc := \text{proc}(i, j, k, l) \frac{(t[k, 1] - t[j, i]) \cdot (t[i, j] - t[l, k])}{(t[i, j] - t[j, i]) \cdot (t[k, l] - t[l, k])} \text{end}$   
 $cc := \text{proc}(i, j, k, l)$   
 $(t[k, l] - t[j, i]) * (t[i, j] - t[l, k]) / ((t[i, j] - t[j, i]) * (t[k, l] - t[l, k]))$   
**end proc**

>  $cc(1, 2, 3, 4)$

$$\frac{(t_{3,4} - t_{2,1}) (t_{1,2} - t_{4,3})}{(t_{1,2} - t_{2,1}) (t_{3,4} - t_{4,3})} \quad (10)$$

>  $\cos(\alpha) :$

>  
>  $C := \text{proc}(i, j, k, l) 2 \cdot \frac{(t[k, 1] - t[j, i]) \cdot (t[i, j] - t[l, k])}{(t[i, j] - t[j, i]) \cdot (t[k, l] - t[l, k])} - 1 \text{end}$   
 $C := \text{proc}(i, j, k, l)$   
 $2 * (t[k, l] - t[j, i]) * (t[i, j] - t[l, k]) / ((t[i, j] - t[j, i]) * (t[k, l] - t[l, k])) - 1$   
**end proc**

>  $C(1, 2, 3, 4)$

$$\frac{2 (t_{3,4} - t_{2,1}) (t_{1,2} - t_{4,3})}{(t_{1,2} - t_{2,1}) (t_{3,4} - t_{4,3})} - 1 \quad (12)$$

> **AMPLITUDE**(12, 13, 24)

>  
>  $Q121324 := -1 + C(1, 2, 1, 3)^2 + C(1, 2, 2, 4)^2 + C(1, 3, 2, 4)^2 + 2 \cdot C(1, 2, 1, 3) \cdot C(1, 2, 4, 2) \cdot C(1, 3, 2, 4)$   
 $Q121324 := -1 + \left( \frac{2 (t_{1,3} - t_{2,1}) (t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1}) (t_{1,3} - t_{3,1})} - 1 \right)^2 + \left( \frac{2 (t_{2,4} - t_{2,1}) (t_{1,2} - t_{4,2})}{(t_{1,2} - t_{2,1}) (t_{2,4} - t_{4,2})} - 1 \right)^2 + \left( \frac{2 (t_{2,4} - t_{3,1}) (t_{1,3} - t_{4,2})}{(t_{1,3} - t_{3,1}) (t_{2,4} - t_{4,2})} - 1 \right)^2 + 2 \left( \frac{2 (t_{1,3} - t_{2,1}) (t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1}) (t_{1,3} - t_{3,1})} - 1 \right) \left( \frac{2 (t_{4,2} - t_{2,1}) (t_{1,2} - t_{2,4})}{(t_{1,2} - t_{2,1}) (t_{4,2} - t_{2,4})} - 1 \right) \left( \frac{2 (t_{2,4} - t_{3,1}) (t_{1,3} - t_{4,2})}{(t_{1,3} - t_{3,1}) (t_{2,4} - t_{4,2})} - 1 \right) \quad (13)$

>  $\text{factor}(\%)$

$$\frac{1}{(t_{1,2} - t_{2,1})^2 (t_{1,3} - t_{3,1})^2 (t_{2,4} - t_{4,2})^2} (4 (t_{1,2} t_{1,3} t_{2,1} - t_{1,2} t_{1,3} t_{3,1} - t_{1,2} t_{2,1} t_{2,4} \\ + t_{1,2} t_{2,1} t_{3,1} - t_{1,2} t_{2,1} t_{4,2} + t_{1,2} t_{2,4} t_{4,2} - t_{1,3} t_{2,1} t_{3,1} + t_{1,3} t_{2,4} t_{3,1} - t_{1,3} t_{2,4} t_{4,2} \\ + t_{1,3} t_{3,1} t_{4,2} + t_{2,1} t_{2,4} t_{4,2} - t_{2,4} t_{3,1} t_{4,2})^2) \quad (14)$$

>

>

>

>  **$Q121324$  in symmetric  $A[i, j] = t[i, j] + t[j, i]$  and antisymmetric  $B[i, j] = t[i, j] + t[j, i]$  coordinates :**

$$\frac{1}{4} \frac{(A_{1,3} - A_{2,4}) (A_{1,2} - A_{2,4}) (A_{1,2} - A_{1,3}) s^3}{B_{1,2} B_{1,3} B_{2,4}} \quad (15)$$

$$-\frac{1}{4} \frac{(A_{1,3} - A_{2,4}) B_{1,2} s}{B_{1,3} B_{2,4}} + \frac{1}{4} \frac{(A_{1,2} - A_{2,4}) B_{1,3} s}{B_{2,4} B_{1,2}} \\ -\frac{1}{4} \frac{(A_{1,2} - A_{1,3}) B_{2,4} s}{B_{1,2} B_{1,3}}$$

$$\begin{aligned}
 > F123 = & \frac{1}{2} \left( \cos\left(\frac{A}{2}\right)^2 + \cos\left(\frac{B}{2}\right)^2 + \cos\left(\frac{C}{2}\right)^2 \right) \\
 - & \frac{1}{8} \frac{B_{1,2}}{B_{2,3}} + \frac{1}{8} \frac{B_{1,2}}{B_{1,3}} + \frac{3}{4} + \frac{1}{8} \frac{B_{1,3}}{B_{2,3}} + \frac{1}{8} \frac{B_{2,3}}{B_{1,3}} \\
 - & \frac{1}{8} \frac{(-A_{2,3} + A_{1,3})^2 s^2}{B_{1,3} B_{2,3}} + \frac{1}{8} \frac{B_{1,3}}{B_{1,2}} - \frac{1}{8} \frac{B_{2,3}}{B_{1,2}} \\
 + & \frac{1}{8} \frac{(A_{1,2} - A_{2,3})^2 s^2}{B_{1,2} B_{2,3}} - \frac{1}{8} \frac{(-A_{1,3} + A_{1,2})^2 s^2}{B_{1,2} B_{1,3}}
 \end{aligned} \tag{16}$$

$$\begin{aligned}
 > G123 = & \left( -1 + \cos(A)^2 + \cos(B)^2 + \cos(C)^2 + 2 \cdot \cos(A) \cdot \cos(B) \cdot \cos(C) \right)^{\frac{1}{2}} \\
 & \frac{1}{8} \frac{(-A_{2,3} + A_{1,3}) B_{1,2} s}{B_{1,3} B_{2,3}} - \frac{1}{8} \frac{(A_{1,2} - A_{2,3}) B_{1,3} s}{B_{1,2} B_{2,3}} \\
 & + \frac{1}{8} \frac{(-A_{1,3} + A_{1,2}) B_{2,3} s}{B_{1,2} B_{1,3}} \\
 & - \frac{1}{8} \frac{(-A_{2,3} + A_{1,3}) (A_{1,2} - A_{2,3}) (-A_{1,3} + A_{1,2}) s^3}{B_{1,2} B_{1,3} B_{2,3}}
 \end{aligned} \tag{17}$$

$$= -1 \left( \frac{2(t_{3,2} - t_{2,1})(t_{1,2} - t_{2,3})}{(t_{1,2} - t_{2,1})(t_{3,2} - t_{2,3})} - 1 \right) \left( \frac{2(t_{2,3} - t_{3,1})(t_{1,3} - t_{3,2})}{(t_{1,3} - t_{3,1})(t_{2,3} - t_{3,2})} - 1 \right)$$

> `factor(%)`

$$\begin{aligned} & \frac{1}{(t_{1,2} - t_{2,1})^2 (t_{1,3} - t_{3,1})^2 (t_{2,3} - t_{3,2})^2} (4(t_{1,2} t_{1,3} t_{2,1} - t_{1,2} t_{1,3} t_{3,1} - t_{1,2} t_{2,1} t_{2,3} \\ & + t_{1,2} t_{2,1} t_{3,1} - t_{1,2} t_{2,1} t_{3,2} + t_{1,2} t_{2,3} t_{3,2} - t_{1,3} t_{2,1} t_{3,1} + t_{1,3} t_{2,3} t_{3,1} - t_{1,3} t_{2,3} t_{3,2} \\ & + t_{1,3} t_{3,1} t_{3,2} + t_{2,1} t_{2,3} t_{3,2} - t_{2,3} t_{3,1} t_{3,2})^2) \end{aligned} \quad (19)$$

>

>

$$\begin{aligned} & > \text{factor}(- (c12^2 - 1) \cdot ((c13^2 - 1)) \cdot (c24^2 - 1) + (c12 \cdot c13 - c23)^2 \cdot (c24^2 - 1) + (c12 \\ & \cdot c24 - c14)^2 \cdot (c13^2 - 1) + (c13 \cdot c24 - c12 \cdot c34)^2 \cdot (c12^2 - 1) + 2 \cdot (c12 \cdot c13 - c23) \\ & \cdot (c12 \cdot c24 - c14) \cdot (c13 \cdot c24 - c12 \cdot c34)) \end{aligned}$$

$$\begin{aligned} & c12^4 c34^2 - 4 c12^3 c13 c24 c34 + 4 c12^2 c13^2 c24^2 + 2 c12^2 c13 c14 c34 \\ & + 2 c12^2 c23 c24 c34 - 4 c12 c13^2 c14 c24 - 4 c12 c13 c23 c24^2 - c12^2 c34^2 \\ & + 2 c12 c13 c24 c34 - 2 c12 c14 c23 c34 + c13^2 c14^2 + 2 c13 c14 c23 c24 + c23^2 c24^2 \\ & + 2 c12 c13 c23 + 2 c12 c14 c24 - c12^2 - c13^2 - c14^2 - c23^2 - c24^2 + 1 \end{aligned} \quad (20)$$

$$\begin{aligned} & > \text{factor}\left(\text{subs}\left(c12 = \frac{\left(x + \frac{1}{x}\right)}{2}, c13 = \frac{\left(y + \frac{1}{y}\right)}{2}, c23 = \frac{\left(z + \frac{1}{z}\right)}{2}, c14 = \frac{\left(u + \frac{1}{u}\right)}{2}, c24 \right. \right. \\ & \left. \left. = \frac{\left(v + \frac{1}{v}\right)}{2}, c34 = \frac{\left(w + \frac{1}{w}\right)}{2}, \% \right)\right): \end{aligned}$$

$$\begin{aligned} & > \text{factor}(- (c12^2 - 1) \cdot (c13^2 - 1) \cdot (c23^2 - 1) + (c12 \cdot c13 - c23)^2 \cdot (c23^2 - 1) + (c12 \cdot c23 \\ & - c13)^2 \cdot (c13^2 - 1) + (c13 \cdot c23 - c12)^2 \cdot (c12^2 - 1) + 2 \cdot (c12 \cdot c13 - c23) \cdot (c12 \cdot c23 \\ & - c13) \cdot (c13 \cdot c23 - c12)) \\ & (-2 c12 c13 c23 + c12^2 + c13^2 + c23^2 - 1)^2 \end{aligned} \quad (21)$$

$$\begin{aligned} & > \text{factor}\left(\text{subs}\left(c12 = \frac{\left(x + \frac{1}{x}\right)}{2}, c13 = \frac{\left(y + \frac{1}{y}\right)}{2}, c23 = \frac{\left(z + \frac{1}{z}\right)}{2}, \% \right)\right) \\ & \frac{1}{16} \frac{(x z - y)^2 (x y - z)^2 (-y z + x)^2 (x y z - 1)^2}{x^4 y^4 z^4} \end{aligned} \quad (22)$$

$$\begin{aligned} & > \text{factor}\left(\text{subs}\left(c12 = \frac{\left(x + \frac{1}{x}\right)}{2}, c13 = \frac{\left(y + \frac{1}{y}\right)}{2}, c23 = \frac{\left(z + \frac{1}{z}\right)}{2}, -1 + c12^2 + c13^2 \right. \right. \\ & \left. \left. + c23^2 + 2 \cdot c12 \cdot c13 \cdot c23 \right)\right) \\ & \frac{1}{4} \frac{(y z + x) (x z + y) (x y z + 1) (x y + z)}{x^2 y^2 z^2} \end{aligned} \quad (23)$$

$$\begin{aligned} & > \text{Sub} := \text{seq}\left(\text{seq}\left(t[i, j] = \frac{(s \cdot A[i, j] + B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right), \text{seq}\left(\text{seq}\left(t[j, i] \right. \right. \\ & \left. \left. = \frac{(s \cdot A[i, j] - B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right): \end{aligned}$$

>

&gt;

$$\begin{aligned}
& > Ev := \frac{1}{16} \cdot ((1 + C(3, 1, 3, 2) + C(2, 3, 2, 4)) \cdot (1 + C(1, 2, 1, 4) + C(4, 1, 4, 3)) + (1 \\
& \quad + C(2, 1, 2, 3) + C(3, 2, 3, 4)) \cdot (1 + C(4, 1, 4, 2) + C(1, 3, 1, 4)) + (1 + C(3, 1, 3, \\
& \quad 2) + C(1, 3, 1, 4)) \cdot (1 + C(2, 1, 2, 4) + C(4, 2, 4, 3)) + (1 + C(1, 2, 1, 3) + C(3, 1, \\
& \quad 3, 4)) \cdot (1 + C(2, 3, 2, 4) + C(4, 1, 4, 2)) + (1 + C(2, 1, 2, 3) + C(1, 2, 1, 4)) \cdot (1 \\
& \quad + C(3, 1, 3, 4) + C(4, 2, 4, 3)) + (1 + C(1, 2, 1, 3) + C(2, 1, 2, 4)) \cdot (1 + C(3, 2, 3, \\
& \quad 4) + C(4, 1, 4, 3))) + \frac{1}{8} \cdot (C(1, 4, 2, 3) \cdot C(1, 3, 2, 4) - C(1, 4, 2, 3) \cdot C(1, 2, 3, 4) \\
& \quad + C(1, 3, 2, 4) \cdot C(1, 2, 3, 4)) \\
& Ev := \frac{1}{16} \left( -1 + \frac{2(t_{3,2} - t_{1,3})(t_{3,1} - t_{2,3})}{(t_{3,1} - t_{1,3})(t_{3,2} - t_{2,3})} + \frac{2(t_{2,4} - t_{3,2})(t_{2,3} - t_{4,2})}{(t_{2,3} - t_{3,2})(t_{2,4} - t_{4,2})} \right) \left( -1 \right) \quad (24) \\
& \quad + \frac{2(t_{1,4} - t_{2,1})(t_{1,2} - t_{4,1})}{(t_{1,2} - t_{2,1})(t_{1,4} - t_{4,1})} + \frac{2(t_{4,3} - t_{1,4})(t_{4,1} - t_{3,4})}{(t_{4,1} - t_{1,4})(t_{4,3} - t_{3,4})} \right) + \frac{1}{16} \left( -1 \right) \\
& \quad + \frac{2(t_{2,3} - t_{1,2})(t_{2,1} - t_{3,2})}{(t_{2,1} - t_{1,2})(t_{2,3} - t_{3,2})} + \frac{2(t_{3,4} - t_{2,3})(t_{3,2} - t_{4,3})}{(t_{3,2} - t_{2,3})(t_{3,4} - t_{4,3})} \right) \left( -1 \right) \\
& \quad + \frac{2(t_{4,2} - t_{1,4})(t_{4,1} - t_{2,4})}{(t_{4,1} - t_{1,4})(t_{4,2} - t_{2,4})} + \frac{2(t_{1,4} - t_{3,1})(t_{1,3} - t_{4,1})}{(t_{1,3} - t_{3,1})(t_{1,4} - t_{4,1})} \right) + \frac{1}{16} \left( -1 \right) \\
& \quad + \frac{2(t_{3,2} - t_{1,3})(t_{3,1} - t_{2,3})}{(t_{3,1} - t_{1,3})(t_{3,2} - t_{2,3})} + \frac{2(t_{1,4} - t_{3,1})(t_{1,3} - t_{4,1})}{(t_{1,3} - t_{3,1})(t_{1,4} - t_{4,1})} \right) \left( -1 \right) \\
& \quad + \frac{2(t_{2,4} - t_{1,2})(t_{2,1} - t_{4,2})}{(t_{2,1} - t_{1,2})(t_{2,4} - t_{4,2})} + \frac{2(t_{4,3} - t_{2,4})(t_{4,2} - t_{3,4})}{(t_{4,2} - t_{2,4})(t_{4,3} - t_{3,4})} \right) + \frac{1}{16} \left( -1 \right) \\
& \quad + \frac{2(t_{1,3} - t_{2,1})(t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1})(t_{1,3} - t_{3,1})} + \frac{2(t_{3,4} - t_{1,3})(t_{3,1} - t_{4,3})}{(t_{3,1} - t_{1,3})(t_{3,4} - t_{4,3})} \right) \left( -1 \right) \\
& \quad + \frac{2(t_{2,4} - t_{3,2})(t_{2,3} - t_{4,2})}{(t_{2,3} - t_{3,2})(t_{2,4} - t_{4,2})} + \frac{2(t_{4,2} - t_{1,4})(t_{4,1} - t_{2,4})}{(t_{4,1} - t_{1,4})(t_{4,2} - t_{2,4})} \right) + \frac{1}{16} \left( -1 \right) \\
& \quad + \frac{2(t_{2,3} - t_{1,2})(t_{2,1} - t_{3,2})}{(t_{2,1} - t_{1,2})(t_{2,3} - t_{3,2})} + \frac{2(t_{1,4} - t_{2,1})(t_{1,2} - t_{4,1})}{(t_{1,2} - t_{2,1})(t_{1,4} - t_{4,1})} \right) \left( -1 \right) \\
& \quad + \frac{2(t_{3,4} - t_{1,3})(t_{3,1} - t_{4,3})}{(t_{3,1} - t_{1,3})(t_{3,4} - t_{4,3})} + \frac{2(t_{4,3} - t_{2,4})(t_{4,2} - t_{3,4})}{(t_{4,2} - t_{2,4})(t_{4,3} - t_{3,4})} \right) + \frac{1}{16} \left( -1 \right) \\
& \quad + \frac{2(t_{1,3} - t_{2,1})(t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1})(t_{1,3} - t_{3,1})} + \frac{2(t_{2,4} - t_{1,2})(t_{2,1} - t_{4,2})}{(t_{2,1} - t_{1,2})(t_{2,4} - t_{4,2})} \right) \left( -1 \right) \\
& \quad + \frac{2(t_{3,4} - t_{2,3})(t_{3,2} - t_{4,3})}{(t_{3,2} - t_{2,3})(t_{3,4} - t_{4,3})} + \frac{2(t_{4,3} - t_{1,4})(t_{4,1} - t_{3,4})}{(t_{4,1} - t_{1,4})(t_{4,3} - t_{3,4})} \right) \\
& \quad + \frac{1}{8} \left( \frac{2(t_{2,3} - t_{4,1})(t_{1,4} - t_{3,2})}{(t_{1,4} - t_{4,1})(t_{2,3} - t_{3,2})} - 1 \right) \left( \frac{2(t_{2,4} - t_{3,1})(t_{1,3} - t_{4,2})}{(t_{1,3} - t_{3,1})(t_{2,4} - t_{4,2})} - 1 \right) \\
& \quad - \frac{1}{8} \left( \frac{2(t_{2,3} - t_{4,1})(t_{1,4} - t_{3,2})}{(t_{1,4} - t_{4,1})(t_{2,3} - t_{3,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{2,1})(t_{1,2} - t_{4,3})}{(t_{1,2} - t_{2,1})(t_{3,4} - t_{4,3})} - 1 \right) \\
& \quad + \frac{1}{8} \left( \frac{2(t_{2,4} - t_{3,1})(t_{1,3} - t_{4,2})}{(t_{1,3} - t_{3,1})(t_{2,4} - t_{4,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{2,1})(t_{1,2} - t_{4,3})}{(t_{1,2} - t_{2,1})(t_{3,4} - t_{4,3})} - 1 \right)
\end{aligned}$$

> **Eodd** := **D4** - **Ev**

> 
$$\begin{aligned} Eodd &:= \text{sort}\left(\text{map}\left(\text{factor}, \text{collect}\left(\text{map}\left(\text{factor}, \text{collect}\left(\text{factor}\left(\text{expand}\left(\text{subs}\left(\text{seq}\left(\text{seq}\left(t[i, j] = \frac{(s \cdot A[i, j] + B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right), \text{seq}\left(\text{seq}\left(t[j, i] = \frac{(s \cdot A[i, j] - B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right)\right), D4 - Ev\right)\right), s\right)\right), [B_{1,2}, B_{1,3}, B_{1,4}, \\ &\quad B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}\right), s\right) : \text{nops}(Eodd) \end{aligned}$$

167 (25)

> **indets**(**Eodd**)

$\{s, A_{1,2}, A_{1,3}, A_{1,4}, A_{2,3}, A_{2,4}, A_{3,4}, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}\}$  (26)

>

> 
$$\begin{aligned} \mathbf{F123} &:= -\frac{1}{8} \frac{B_{1,2}}{B_{2,3}} + \frac{1}{8} \frac{B_{1,2}}{B_{1,3}} + \frac{3}{4} + \frac{1}{8} \frac{B_{1,3}}{B_{2,3}} + \frac{1}{8} \frac{B_{2,3}}{B_{1,3}} - \frac{1}{8} \frac{(-A_{2,3} + A_{1,3})^2 s^2}{B_{2,3} B_{1,3}} \\ &\quad + \frac{1}{8} \frac{B_{1,3}}{B_{1,2}} - \frac{1}{8} \frac{B_{2,3}}{B_{1,2}} + \frac{1}{8} \frac{(A_{1,2} - A_{2,3})^2 s^2}{B_{1,2} B_{2,3}} - \frac{1}{8} \frac{(-A_{1,3} + A_{1,2})^2 s^2}{B_{1,2} B_{1,3}} : \\ \mathbf{F124} &:= -\frac{1}{8} \frac{B_{1,2}}{B_{2,4}} + \frac{1}{8} \frac{B_{1,2}}{B_{1,4}} + \frac{3}{4} + \frac{1}{8} \frac{B_{1,4}}{B_{2,4}} + \frac{1}{8} \frac{B_{2,4}}{B_{1,4}} - \frac{1}{8} \frac{(-A_{2,4} + A_{1,4})^2 s^2}{B_{2,4} B_{1,4}} \\ &\quad + \frac{1}{8} \frac{B_{1,4}}{B_{1,2}} - \frac{1}{8} \frac{B_{2,4}}{B_{1,2}} + \frac{1}{8} \frac{(A_{1,2} - A_{2,4})^2 s^2}{B_{1,2} B_{2,4}} - \frac{1}{8} \frac{(-A_{1,4} + A_{1,2})^2 s^2}{B_{1,2} B_{1,4}} \end{aligned}$$

>

> 
$$\begin{aligned} \mathbf{F134} &:= -\frac{1}{8} \frac{B_{1,3}}{B_{3,4}} + \frac{1}{8} \frac{B_{1,3}}{B_{1,4}} + \frac{3}{4} + \frac{1}{8} \frac{B_{1,4}}{B_{3,4}} + \frac{1}{8} \frac{B_{3,4}}{B_{1,4}} - \frac{1}{8} \frac{(-A_{3,4} + A_{1,4})^2 s^2}{B_{3,4} B_{1,4}} \\ &\quad + \frac{1}{8} \frac{B_{1,4}}{B_{1,3}} - \frac{1}{8} \frac{B_{3,4}}{B_{1,3}} + \frac{1}{8} \frac{(A_{1,3} - A_{3,4})^2 s^2}{B_{1,3} B_{3,4}} - \frac{1}{8} \frac{(-A_{1,4} + A_{1,3})^2 s^2}{B_{1,3} B_{1,4}} : \\ \mathbf{F234} &:= -\frac{1}{8} \frac{B_{2,3}}{B_{3,4}} + \frac{1}{8} \frac{B_{2,3}}{B_{2,4}} + \frac{3}{4} + \frac{1}{8} \frac{B_{2,4}}{B_{3,4}} + \frac{1}{8} \frac{B_{3,4}}{B_{2,4}} - \frac{1}{8} \frac{(-A_{3,4} + A_{2,4})^2 s^2}{B_{3,4} B_{2,4}} \\ &\quad + \frac{1}{8} \frac{B_{2,4}}{B_{2,3}} - \frac{1}{8} \frac{B_{3,4}}{B_{2,3}} + \frac{1}{8} \frac{(A_{2,3} - A_{3,4})^2 s^2}{B_{2,3} B_{3,4}} - \frac{1}{8} \frac{(-A_{2,4} + A_{2,3})^2 s^2}{B_{2,3} B_{2,4}} : \end{aligned}$$

> 
$$\begin{aligned} \mathbf{G123} &:= \text{map}\left(\text{factor}, \text{collect}\left(\text{factor}\left(\text{expand}\left(\text{subs}\left(\text{seq}\left(\text{seq}\left(t[i, j] = \frac{(s \cdot A[i, j] + B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 3\right), \text{seq}\left(\text{seq}\left(t[j, i] = \frac{(s \cdot A[i, j] - B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 3\right)\right), \frac{1}{8} \frac{(-A_{2,3} + A_{1,3}) B_{1,2} s}{B_{2,3} B_{1,3}} - \frac{1}{8} \frac{(A_{1,2} - A_{2,3}) B_{1,3} s}{B_{1,2} B_{2,3}} + \frac{1}{8} \frac{(-A_{1,3} + A_{1,2}) B_{2,3} s}{B_{1,2} B_{1,3}} \right. \\ &\quad \left. - \frac{1}{8} \frac{(-A_{2,3} + A_{1,3}) (A_{1,2} - A_{2,3}) (-A_{1,3} + A_{1,2}) s^3}{B_{1,2} B_{1,3} B_{2,3}} \right)\right), [B_{1,2}, B_{1,3}, B_{2,3}], \text{distributed}\right) \end{aligned}$$

$$G123 := \frac{1}{8} \frac{(A_{1,3} - A_{2,3}) B_{1,2} s}{B_{2,3} B_{1,3}} - \frac{1}{8} \frac{(A_{1,2} - A_{2,3}) B_{1,3} s}{B_{2,3} B_{1,2}} + \frac{1}{8} \frac{(A_{1,2} - A_{1,3}) B_{2,3} s}{B_{1,2} B_{1,3}}$$
 (27)

$$-\frac{1}{8} \frac{(A_{1,3} - A_{2,3})(A_{1,2} - A_{2,3})(A_{1,2} - A_{1,3}) s^3}{B_{1,2} B_{2,3} B_{1,3}}$$

$$\begin{aligned} > \mathbf{G124} := & \frac{1}{8} \frac{(-A_{2,4} + A_{1,4}) B_{1,2} s}{B_{2,4} B_{1,4}} - \frac{1}{8} \frac{(A_{1,2} - A_{2,4}) B_{1,4} s}{B_{1,2} B_{2,4}} + \frac{1}{8} \frac{(-A_{1,4} + A_{1,2}) B_{2,4} s}{B_{1,2} B_{1,4}} \\ & - \frac{1}{8} \frac{(-A_{2,4} + A_{1,4})(A_{1,2} - A_{2,4})(-A_{1,4} + A_{1,2}) s^3}{B_{1,2} B_{1,4} B_{2,4}} : \end{aligned}$$

$$\begin{aligned} > \mathbf{G134} := & \frac{1}{8} \frac{(-A_{3,4} + A_{1,4}) B_{1,3} s}{B_{3,4} B_{1,4}} - \frac{1}{8} \frac{(A_{1,3} - A_{3,4}) B_{1,4} s}{B_{1,3} B_{3,4}} + \frac{1}{8} \frac{(-A_{1,4} + A_{1,3}) B_{3,4} s}{B_{1,3} B_{1,4}} \\ & - \frac{1}{8} \frac{(-A_{3,4} + A_{1,4})(A_{1,3} - A_{3,4})(-A_{1,4} + A_{1,3}) s^3}{B_{1,3} B_{1,4} B_{3,4}} : \end{aligned}$$

$$\begin{aligned} > \mathbf{G234} := & \frac{1}{8} \frac{(-A_{3,4} + A_{2,4}) B_{2,3} s}{B_{3,4} B_{2,4}} - \frac{1}{8} \frac{(A_{2,3} - A_{3,4}) B_{2,4} s}{B_{2,3} B_{3,4}} + \frac{1}{8} \frac{(-A_{2,4} + A_{2,3}) B_{3,4} s}{B_{2,3} B_{2,4}} \\ & - \frac{1}{8} \frac{(-A_{3,4} + A_{2,4})(A_{2,3} - A_{3,4})(-A_{2,4} + A_{2,3}) s^3}{B_{2,3} B_{2,4} B_{3,4}} : \end{aligned}$$

$$> \mathbf{Eodd1} := \text{sort}\left(Eodd + \frac{1}{2} \cdot (G123 + G124 + G134 + G234), s\right) : \text{nops}(Eodd1)$$

151 (28)

$$\begin{aligned} > g121324 := & -\frac{1}{32} \frac{(A_{1,3} - A_{2,4}) B_{1,2} s}{B_{1,3} B_{2,4}} + \frac{1}{32} \frac{(A_{1,2} - A_{2,4}) B_{1,3} s}{B_{1,2} B_{2,4}} \\ & - \frac{1}{32} \frac{(-A_{1,3} + A_{1,2}) B_{2,4} s}{B_{1,2} B_{1,3}} + \frac{1}{32} \frac{(A_{1,3} - A_{2,4})(A_{1,2} - A_{2,4})(-A_{1,3} + A_{1,2}) s^3}{B_{1,2} B_{1,3} B_{2,4}} : \end{aligned}$$

$$\begin{aligned} > g121334 := & \frac{1}{32} \frac{(A_{1,3} - A_{3,4}) B_{1,2} s}{B_{1,3} B_{3,4}} - \frac{1}{32} \frac{(-A_{3,4} + A_{1,2}) B_{1,3} s}{B_{1,2} B_{3,4}} \\ & + \frac{1}{32} \frac{(-A_{1,3} + A_{1,2}) B_{3,4} s}{B_{1,2} B_{1,3}} - \frac{1}{32} \frac{(A_{1,3} - A_{3,4})(-A_{3,4} + A_{1,2})(-A_{1,3} + A_{1,2}) s^3}{B_{1,2} B_{1,3} B_{3,4}} : \end{aligned}$$

$$\begin{aligned} > g121423 := & -\frac{1}{32} \frac{(A_{1,4} - A_{2,3}) B_{1,2} s}{B_{1,4} B_{2,3}} + \frac{1}{32} \frac{(A_{1,2} - A_{2,3}) B_{1,4} s}{B_{1,2} B_{2,3}} - \frac{1}{32} \frac{(A_{1,2} - A_{1,4}) B_{2,3} s}{B_{1,2} B_{1,4}} \\ & + \frac{1}{32} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{2,3})(A_{1,2} - A_{1,4}) s^3}{B_{1,2} B_{1,4} B_{2,3}} : \end{aligned}$$

$$\begin{aligned} > g121434 := & -\frac{1}{32} \frac{(A_{1,4} - A_{3,4}) B_{1,2} s}{B_{1,4} B_{3,4}} + \frac{1}{32} \frac{(-A_{3,4} + A_{1,2}) B_{1,4} s}{B_{1,2} B_{3,4}} \\ & - \frac{1}{32} \frac{(A_{1,2} - A_{1,4}) B_{3,4} s}{B_{1,2} B_{1,4}} + \frac{1}{32} \frac{(A_{1,4} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{1,4}) s^3}{B_{1,2} B_{1,4} B_{3,4}} : \end{aligned}$$

$$\begin{aligned} > g122334 := & -\frac{1}{32} \frac{(A_{2,3} - A_{3,4}) B_{1,2} s}{B_{2,3} B_{3,4}} + \frac{1}{32} \frac{(-A_{3,4} + A_{1,2}) B_{2,3} s}{B_{1,2} B_{3,4}} \\ & - \frac{1}{32} \frac{(A_{1,2} - A_{2,3}) B_{3,4} s}{B_{1,2} B_{2,3}} + \frac{1}{32} \frac{(A_{2,3} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{2,3}) s^3}{B_{1,2} B_{2,3} B_{3,4}} : \end{aligned}$$

$$> g122434 := +\frac{1}{32} \frac{(A_{2,4} - A_{3,4}) B_{1,2} s}{B_{2,4} B_{3,4}} - \frac{1}{32} \frac{(-A_{3,4} + A_{1,2}) B_{2,4} s}{B_{1,2} B_{3,4}}$$

$$\begin{aligned}
& + \frac{1}{32} \frac{(A_{1,2} - A_{2,4}) B_{3,4} s}{B_{1,2} B_{2,4}} - \frac{1}{32} \frac{(A_{2,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,4}) s^3}{B_{1,2} B_{2,4} B_{3,4}} : \\
> g131423 & := + \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) B_{1,3} s}{B_{1,4} B_{2,3}} - \frac{1}{32} \frac{(-A_{2,3} + A_{1,3}) B_{1,4} s}{B_{1,3} B_{2,3}} + \frac{1}{32} \frac{(A_{1,3} - A_{1,4}) B_{2,3} s}{B_{1,3} B_{1,4}} \\
& - \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) (-A_{2,3} + A_{1,3}) (A_{1,3} - A_{1,4}) s^3}{B_{1,3} B_{1,4} B_{2,3}} : \\
> g131424 & := - \frac{1}{32} \frac{(A_{1,4} - A_{2,4}) B_{1,3} s}{B_{1,4} B_{2,4}} + \frac{1}{32} \frac{(A_{1,3} - A_{2,4}) B_{1,4} s}{B_{1,3} B_{2,4}} \\
& - \frac{1}{32} \frac{(A_{1,3} - A_{1,4}) B_{2,4} s}{B_{1,3} B_{1,4}} + \frac{1}{32} \frac{(A_{1,4} - A_{2,4}) (A_{1,3} - A_{2,4}) (A_{1,3} - A_{1,4}) s^3}{B_{1,3} B_{1,4} B_{2,4}} : \\
> g132324 & := + \frac{1}{32} \frac{(A_{2,3} - A_{2,4}) B_{1,3} s}{B_{2,3} B_{2,4}} - \frac{1}{32} \frac{(A_{1,3} - A_{2,4}) B_{2,3} s}{B_{1,3} B_{2,4}} + \frac{1}{32} \frac{(-A_{2,3} + A_{1,3}) B_{2,4} s}{B_{1,3} B_{2,3}} \\
& - \frac{1}{32} \frac{(A_{2,3} - A_{2,4}) (A_{1,3} - A_{2,4}) (-A_{2,3} + A_{1,3}) s^3}{B_{1,3} B_{2,3} B_{2,4}} : \\
> g132434 & := - \frac{1}{32} \frac{(A_{2,4} - A_{3,4}) B_{1,3} s}{B_{2,4} B_{3,4}} + \frac{1}{32} \frac{(A_{1,3} - A_{3,4}) B_{2,4} s}{B_{1,3} B_{3,4}} \\
& - \frac{1}{32} \frac{(A_{1,3} - A_{2,4}) B_{3,4} s}{B_{1,3} B_{2,4}} + \frac{1}{32} \frac{(A_{2,4} - A_{3,4}) (A_{1,3} - A_{3,4}) (A_{1,3} - A_{2,4}) s^3}{B_{1,3} B_{2,4} B_{3,4}} : \\
> g142324 & := - \frac{1}{32} \frac{(A_{2,3} - A_{2,4}) B_{1,4} s}{B_{2,3} B_{2,4}} + \frac{1}{32} \frac{(A_{1,4} - A_{2,4}) B_{2,3} s}{B_{1,4} B_{2,4}} - \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) B_{2,4} s}{B_{1,4} B_{2,3}} \\
& + \frac{1}{32} \frac{(A_{2,3} - A_{2,4}) (A_{1,4} - A_{2,4}) (A_{1,4} - A_{2,3}) s^3}{B_{1,4} B_{2,3} B_{2,4}} : \\
> g142334 & := \frac{1}{32} \frac{(A_{2,3} - A_{3,4}) B_{1,4} s}{B_{2,3} B_{3,4}} - \frac{1}{32} \frac{(A_{1,4} - A_{3,4}) B_{2,3} s}{B_{1,4} B_{3,4}} + \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) B_{3,4} s}{B_{1,4} B_{2,3}} \\
& - \frac{1}{32} \frac{(A_{2,3} - A_{3,4}) (A_{1,4} - A_{3,4}) (A_{1,4} - A_{2,3}) s^3}{B_{1,4} B_{2,3} B_{3,4}} :
\end{aligned}$$

$$\begin{aligned}
> Eodd2 & := sort(Eodd1 - g121324 - g121334 - g121423 - g121434 - g122334 \\
& - g122434 - g131423 - g131424 - g132324 - g132434 - g142324 - g142334, s) : \\
nops(\%) &
\end{aligned}$$

$$g121324c1423 := map(factor, collect( g121324·subs(Sub, C(1,4,2,3)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], distributed) )$$

$$\begin{aligned}
g121324c1423 & := - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (A_{1,2} - A_{2,4}) (A_{1,2} - A_{1,3}) (A_{1,4} - A_{2,3})^2 s^5}{B_{2,3} B_{1,4} B_{2,4} B_{1,2} B_{1,3}} \quad (29) \\
& + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (A_{1,4} - A_{2,3})^2 s^3 B_{1,2}}{B_{2,3} B_{1,4} B_{1,3} B_{2,4}} - \frac{1}{64} \frac{(A_{1,2} - A_{2,4}) (A_{1,4} - A_{2,3})^2 s^3 B_{1,3}}{B_{2,3} B_{1,4} B_{2,4} B_{1,2}} \\
& + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (A_{1,2} - A_{2,4}) (A_{1,2} - A_{1,3}) s^3 B_{1,4}}{B_{2,3} B_{2,4} B_{1,2} B_{1,3}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{1,3}) (A_{1,4} - A_{2,3})^2 B_{2,4} s^3}{B_{2,3} B_{1,2} B_{1,3} B_{1,4}}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3})(A_{1,2} - A_{2,4})(A_{1,2} - A_{1,3})B_{2,3}s^3}{B_{2,4}B_{1,2}B_{1,3}B_{1,4}} \\
& - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3})B_{1,2}B_{1,4}s}{B_{2,3}B_{1,3}B_{2,4}} - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3})B_{2,3}B_{1,2}s}{B_{1,3}B_{1,4}B_{2,4}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{2,4})sB_{1,3}B_{1,4}}{B_{2,3}B_{2,4}B_{1,2}} + \frac{1}{64} \frac{(A_{1,2} - A_{2,4})B_{2,3}sB_{1,3}}{B_{2,4}B_{1,2}B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{1,3})B_{2,4}sB_{1,4}}{B_{2,3}B_{1,2}B_{1,3}} - \frac{1}{64} \frac{(A_{1,2} - A_{1,3})B_{2,3}B_{2,4}s}{B_{1,2}B_{1,3}B_{1,4}}
\end{aligned}$$

>  $g121334c1423 := \text{map}(\text{factor}, \text{collect}(-g121334 \cdot \text{subs}(\text{Sub}, C(1, 4, 2, 3)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g121334c1423 := & -\frac{1}{64} \frac{(A_{1,3} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{1,3})(A_{1,4} - A_{2,3})^2 s^5}{B_{2,3}B_{1,4}B_{3,4}B_{1,3}B_{1,2}} \quad (30) \\
& + \frac{1}{64} \frac{(A_{1,3} - A_{3,4})(A_{1,4} - A_{2,3})^2 s^3 B_{1,2}}{B_{2,3}B_{1,4}B_{1,3}B_{3,4}} - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2})(A_{1,4} - A_{2,3})^2 s^3 B_{1,3}}{B_{2,3}B_{1,4}B_{3,4}B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{1,3} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{1,3})s^3 B_{1,4}}{B_{2,3}B_{3,4}B_{1,2}B_{1,3}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{1,3})(A_{1,4} - A_{2,3})^2 B_{3,4}s^3}{B_{2,3}B_{1,2}B_{1,3}B_{1,4}} \\
& + \frac{1}{64} \frac{(A_{1,3} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{1,3})B_{2,3}s^3}{B_{3,4}B_{1,2}B_{1,3}B_{1,4}} - \frac{1}{64} \frac{(A_{1,3} - A_{3,4})sB_{1,2}B_{1,4}}{B_{2,3}B_{3,4}B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{3,4})B_{2,3}sB_{1,2}}{B_{3,4}B_{1,3}B_{1,4}} + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2})B_{1,3}B_{1,4}s}{B_{1,2}B_{2,3}B_{3,4}} \\
& + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2})B_{1,3}B_{2,3}s}{B_{1,2}B_{1,4}B_{3,4}} - \frac{1}{64} \frac{(A_{1,2} - A_{1,3})B_{3,4}sB_{1,4}}{B_{2,3}B_{1,2}B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{1,3})B_{2,3}B_{3,4}s}{B_{1,2}B_{1,3}B_{1,4}}
\end{aligned}$$

>

>  $g121423c1324 := \text{map}(\text{factor}, \text{collect}(g121423 \cdot \text{subs}(\text{Sub}, C(1, 3, 2, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g121423c1324 := & -\frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{2,3})(A_{1,2} - A_{1,4})(-A_{2,4} + A_{1,3})^2 s^5}{B_{2,3}B_{1,4}B_{2,4}B_{1,2}B_{1,3}} \quad (31) \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(-A_{2,4} + A_{1,3})^2 s^3 B_{1,2}}{B_{2,3}B_{1,4}B_{1,3}B_{2,4}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{2,3})(A_{1,2} - A_{1,4})s^3 B_{1,3}}{B_{2,3}B_{1,4}B_{2,4}B_{1,2}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{2,3})(-A_{2,4} + A_{1,3})^2 s^3 B_{1,4}}{B_{2,3}B_{2,4}B_{1,2}B_{1,3}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{2,3})(A_{1,2} - A_{1,4})B_{2,4}s^3}{B_{2,3}B_{1,2}B_{1,3}B_{1,4}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{1,4})(-A_{2,4} + A_{1,3})^2 B_{2,3}s^3}{B_{2,4}B_{1,2}B_{1,3}B_{1,4}} - \frac{1}{64} \frac{(A_{1,4} - A_{2,3})B_{1,3}B_{1,2}s}{B_{2,3}B_{1,4}B_{2,4}}
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) B_{1,2} B_{2,4} s}{B_{1,3} B_{1,4} B_{2,3}} + \frac{1}{64} \frac{(A_{1,2} - A_{2,3}) s B_{1,3} B_{1,4}}{B_{2,3} B_{2,4} B_{1,2}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{1,4}) B_{2,3} s B_{1,3}}{B_{2,4} B_{1,2} B_{1,4}} + \frac{1}{64} \frac{(A_{1,2} - A_{2,3}) B_{2,4} s B_{1,4}}{B_{2,3} B_{1,2} B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{1,4}) B_{2,3} B_{2,4} s}{B_{1,2} B_{1,3} B_{1,4}}
\end{aligned}$$

>  $g121434c1324 := \text{map}(\text{factor}, \text{collect}(-g121434 \cdot \text{subs}(\text{Sub}, C(1, 3, 2, 4)), [s, B_{1,2}, B_{1,3},$   
 $B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g121434c1324 := & \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{1,4}) (-A_{2,4} + A_{1,3})^2 s^5}{B_{2,4} B_{1,3} B_{3,4} B_{1,4} B_{1,2}} \quad (32) \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) (-A_{2,4} + A_{1,3})^2 s^3 B_{1,2}}{B_{2,4} B_{1,3} B_{1,4} B_{3,4}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{1,4}) s^3 B_{1,3}}{B_{2,4} B_{3,4} B_{1,4} B_{1,2}} \\
& + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) (-A_{2,4} + A_{1,3})^2 s^3 B_{1,4}}{B_{2,4} B_{3,4} B_{1,2} B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{1,4}) (-A_{2,4} + A_{1,3})^2 B_{3,4} s^3}{B_{2,4} B_{1,2} B_{1,3} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{1,4}) B_{2,4} s^3}{B_{3,4} B_{1,2} B_{1,3} B_{1,4}} + \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) s B_{1,2} B_{1,3}}{B_{2,4} B_{3,4} B_{1,4}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{3,4}) B_{2,4} s B_{1,2}}{B_{3,4} B_{1,3} B_{1,4}} - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{1,3} B_{1,4} s}{B_{2,4} B_{3,4} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{1,4}) B_{3,4} s B_{1,3}}{B_{2,4} B_{1,2} B_{1,4}} - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{1,4} B_{2,4} s}{B_{3,4} B_{1,3} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{1,4}) B_{2,4} B_{3,4} s}{B_{1,2} B_{1,3} B_{1,4}}
\end{aligned}$$

>  $g122334c1324 := \text{map}(\text{factor}, \text{collect}(-g122334 \cdot \text{subs}(\text{Sub}, C(1, 3, 2, 4)), [s, B_{1,2}, B_{1,3},$   
 $B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g122334c1324 := & \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,3}) (-A_{2,4} + A_{1,3})^2 s^5}{B_{2,4} B_{1,3} B_{3,4} B_{2,3} B_{1,2}} \quad (33) \\
& - \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) (-A_{2,4} + A_{1,3})^2 s^3 B_{1,2}}{B_{2,4} B_{1,3} B_{2,3} B_{3,4}} \\
& - \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,3}) s^3 B_{1,3}}{B_{2,4} B_{3,4} B_{2,3} B_{1,2}} \\
& + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) (-A_{2,4} + A_{1,3})^2 s^3 B_{2,3}}{B_{2,4} B_{3,4} B_{1,2} B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{2,3}) (-A_{2,4} + A_{1,3})^2 B_{3,4} s^3}{B_{2,3} B_{2,4} B_{1,2} B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,3}) B_{2,4} s^3}{B_{3,4} B_{1,2} B_{1,3} B_{2,3}} + \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) s B_{1,2} B_{1,3}}{B_{2,4} B_{3,4} B_{2,3}}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{64} \frac{(A_{2,3} - A_{3,4}) B_{2,4} s B_{1,2}}{B_{2,3} B_{3,4} B_{1,3}} - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{1,3} B_{2,3} s}{B_{2,4} B_{3,4} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{2,3}) B_{3,4} s B_{1,3}}{B_{2,3} B_{2,4} B_{1,2}} - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{2,3} B_{2,4} s}{B_{3,4} B_{1,3} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{1,2} - A_{2,3}) B_{2,4} B_{3,4} s}{B_{2,3} B_{1,2} B_{1,3}}
\end{aligned}$$

>  $g122434c1423 := \text{map}(\text{factor}, \text{collect}(-g122434 \cdot \text{subs}(Sub, C(1, 4, 2, 3)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g122434c1423 := & - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,4}) (A_{1,4} - A_{2,3})^2 s^5}{B_{2,3} B_{1,4} B_{3,4} B_{2,4} B_{1,2}} \quad (34) \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (A_{1,4} - A_{2,3})^2 s^3 B_{1,2}}{B_{2,3} B_{1,4} B_{2,4} B_{3,4}} \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,4}) s^3 B_{1,4}}{B_{2,4} B_{3,4} B_{2,3} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (-A_{3,4} + A_{1,2}) (A_{1,2} - A_{2,4}) s^3 B_{2,3}}{B_{2,4} B_{3,4} B_{1,2} B_{1,4}} \\
& - \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) (A_{1,4} - A_{2,3})^2 s^3 B_{2,4}}{B_{2,3} B_{1,4} B_{3,4} B_{1,2}} + \frac{1}{64} \frac{(A_{1,2} - A_{2,4}) (A_{1,4} - A_{2,3})^2 B_{3,4} s^3}{B_{2,3} B_{1,4} B_{2,4} B_{1,2}} \\
& - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) s B_{1,2} B_{1,4}}{B_{2,4} B_{3,4} B_{2,3}} - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) B_{2,3} s B_{1,2}}{B_{2,4} B_{3,4} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{2,4}) B_{3,4} s B_{1,4}}{B_{2,3} B_{2,4} B_{1,2}} + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{2,4} B_{1,4} s}{B_{1,2} B_{2,3} B_{3,4}} \\
& - \frac{1}{64} \frac{(A_{1,2} - A_{2,4}) B_{3,4} s B_{2,3}}{B_{2,4} B_{1,2} B_{1,4}} + \frac{1}{64} \frac{(-A_{3,4} + A_{1,2}) B_{2,3} B_{2,4} s}{B_{1,2} B_{1,4} B_{3,4}}
\end{aligned}$$

>

>  $g131423c1234 := \text{map}(\text{factor}, \text{collect}(g131423 \cdot \text{subs}(Sub, C(1, 2, 3, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g131423c1234 := & \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) (A_{1,3} - A_{2,3}) (A_{1,3} - A_{1,4}) (-A_{3,4} + A_{1,2})^2 s^5}{B_{2,3} B_{1,4} B_{3,4} B_{1,3} B_{1,2}} \quad (35) \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) (A_{1,3} - A_{2,3}) (A_{1,3} - A_{1,4}) s^3 B_{1,2}}{B_{2,3} B_{1,4} B_{1,3} B_{3,4}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,3}}{B_{2,3} B_{1,4} B_{3,4} B_{1,2}} + \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,4}}{B_{2,3} B_{3,4} B_{1,2} B_{1,3}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) (A_{1,3} - A_{2,3}) (A_{1,3} - A_{1,4}) B_{3,4} s^3}{B_{2,3} B_{1,2} B_{1,3} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) (-A_{3,4} + A_{1,2})^2 B_{2,3} s^3}{B_{3,4} B_{1,2} B_{1,3} B_{1,4}} + \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) B_{1,3} B_{1,2} s}{B_{2,3} B_{1,4} B_{3,4}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) s B_{1,2} B_{1,4}}{B_{2,3} B_{3,4} B_{1,3}} + \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) B_{2,3} s B_{1,2}}{B_{3,4} B_{1,3} B_{1,4}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) B_{3,4} B_{1,3} s}{B_{2,3} B_{1,2} B_{1,4}} - \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) B_{3,4} s B_{1,4}}{B_{2,3} B_{1,2} B_{1,3}}
\end{aligned}$$

$$+ \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) B_{2,3} B_{3,4} s}{B_{1,2} B_{1,3} B_{1,4}}$$

>

>  $g131424c1234 := \text{map}(\text{factor}, \text{collect}(-g131424\cdot\text{subs}(\text{Sub}, C(1, 2, 3, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned} g131424c1234 := & \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{1,4}) (-A_{3,4} + A_{1,2})^2 s^5}{B_{2,4} B_{1,3} B_{3,4} B_{1,4} B_{1,2}} \\ & - \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{1,4}) s^3 B_{1,2}}{B_{2,4} B_{1,3} B_{1,4} B_{3,4}} \\ & - \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,3}}{B_{2,4} B_{3,4} B_{1,4} B_{1,2}} \\ & + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,4}}{B_{2,4} B_{3,4} B_{1,2} B_{1,3}} \\ & - \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{1,4}) B_{3,4} s^3}{B_{2,4} B_{1,2} B_{1,3} B_{1,4}} \\ & - \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) (-A_{3,4} + A_{1,2})^2 B_{2,4} s^3}{B_{3,4} B_{1,2} B_{1,3} B_{1,4}} + \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) s B_{1,2} B_{1,3}}{B_{2,4} B_{3,4} B_{1,4}} \\ & - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{1,2} B_{1,4} s}{B_{2,4} B_{3,4} B_{1,3}} + \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) B_{2,4} s B_{1,2}}{B_{3,4} B_{1,3} B_{1,4}} \\ & + \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) B_{3,4} s B_{1,3}}{B_{2,4} B_{1,2} B_{1,4}} - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{3,4} B_{1,4} s}{B_{1,2} B_{1,3} B_{2,4}} \\ & + \frac{1}{64} \frac{(A_{1,3} - A_{1,4}) B_{2,4} B_{3,4} s}{B_{1,2} B_{1,3} B_{1,4}} \end{aligned} \quad (36)$$

>

>  $g132324c1234 := \text{map}(\text{factor}, \text{collect}(-g132324\cdot\text{subs}(\text{Sub}, C(1, 2, 3, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned} g132324c1234 := & - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 s^5}{B_{2,4} B_{1,3} B_{3,4} B_{2,3} B_{1,2}} \\ & + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{2,3}) s^3 B_{1,2}}{B_{2,4} B_{1,3} B_{2,3} B_{3,4}} \\ & + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,3}}{B_{2,4} B_{3,4} B_{2,3} B_{1,2}} \\ & - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (-A_{3,4} + A_{1,2})^2 s^3 B_{2,3}}{B_{2,4} B_{3,4} B_{1,2} B_{1,3}} \\ & + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (-A_{2,4} + A_{1,3}) (A_{1,3} - A_{2,3}) B_{3,4} s^3}{B_{2,4} B_{1,2} B_{1,3} B_{2,3}} \\ & + \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 B_{2,4} s^3}{B_{2,3} B_{3,4} B_{1,2} B_{1,3}} - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) s B_{1,2} B_{1,3}}{B_{2,4} B_{3,4} B_{2,3}} \\ & + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{1,2} B_{2,3} s}{B_{1,3} B_{2,4} B_{3,4}} - \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) B_{2,4} s B_{1,2}}{B_{2,3} B_{3,4} B_{1,3}} \end{aligned} \quad (37)$$

$$\begin{aligned}
& - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) B_{3,4} s B_{1,3}}{B_{2,3} B_{2,4} B_{1,2}} + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{3,4} B_{2,3} s}{B_{1,3} B_{1,2} B_{2,4}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{2,3}) B_{2,4} B_{3,4} s}{B_{2,3} B_{1,2} B_{1,3}}
\end{aligned}$$

>  $g132434c1423 := \text{map}(\text{factor}, \text{collect}(\text{g132434}\cdot\text{subs}(\text{Sub}, C(1, 4, 2, 3)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g132434c1423 := & - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (A_{1,3} - A_{3,4}) (-A_{2,4} + A_{1,3}) (A_{1,4} - A_{2,3})^2 s^5}{B_{2,3} B_{1,4} B_{3,4} B_{2,4} B_{1,3}} \quad (38) \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (A_{1,4} - A_{2,3})^2 s^3 B_{1,3}}{B_{2,3} B_{1,4} B_{2,4} B_{3,4}} \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (A_{1,3} - A_{3,4}) (-A_{2,4} + A_{1,3}) s^3 B_{1,4}}{B_{2,4} B_{3,4} B_{2,3} B_{1,3}} \\
& + \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) (A_{1,3} - A_{3,4}) (-A_{2,4} + A_{1,3}) s^3 B_{2,3}}{B_{2,4} B_{3,4} B_{1,3} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{3,4}) (A_{1,4} - A_{2,3})^2 s^3 B_{2,4}}{B_{2,3} B_{1,4} B_{1,3} B_{3,4}} + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) (A_{1,4} - A_{2,3})^2 B_{3,4} s^3}{B_{2,3} B_{1,4} B_{1,3} B_{2,4}} \\
& - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) s B_{1,3} B_{1,4}}{B_{2,4} B_{3,4} B_{2,3}} - \frac{1}{64} \frac{(A_{2,4} - A_{3,4}) B_{2,3} s B_{1,3}}{B_{2,4} B_{3,4} B_{1,4}} \\
& - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{3,4} B_{1,4} s}{B_{1,3} B_{2,3} B_{2,4}} + \frac{1}{64} \frac{(A_{1,3} - A_{3,4}) B_{2,4} s B_{1,4}}{B_{2,3} B_{3,4} B_{1,3}} \\
& - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3}) B_{3,4} B_{2,3} s}{B_{1,3} B_{1,4} B_{2,4}} + \frac{1}{64} \frac{(A_{1,3} - A_{3,4}) B_{2,4} s B_{2,3}}{B_{3,4} B_{1,3} B_{1,4}}
\end{aligned}$$

>  $g142324c1234 := \text{map}(\text{factor}, \text{collect}(\text{g142324}\cdot\text{subs}(\text{Sub}, C(1, 2, 3, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$$\begin{aligned}
g142324c1234 := & - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (A_{1,4} - A_{2,4}) (A_{1,4} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 s^5}{B_{2,3} B_{1,4} B_{3,4} B_{2,4} B_{1,2}} \quad (39) \\
& + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (A_{1,4} - A_{2,4}) (A_{1,4} - A_{2,3}) s^3 B_{1,2}}{B_{2,3} B_{1,4} B_{2,4} B_{3,4}} \\
& + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (-A_{3,4} + A_{1,2})^2 s^3 B_{1,4}}{B_{2,4} B_{3,4} B_{2,3} B_{1,2}} - \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) (-A_{3,4} + A_{1,2})^2 s^3 B_{2,3}}{B_{2,4} B_{3,4} B_{1,4} B_{1,2}} \\
& + \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) (A_{1,4} - A_{2,4}) (A_{1,4} - A_{2,3}) B_{3,4} s^3}{B_{2,4} B_{1,4} B_{1,2} B_{2,3}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) (-A_{3,4} + A_{1,2})^2 B_{2,4} s^3}{B_{2,3} B_{1,4} B_{3,4} B_{1,2}} - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) s B_{1,2} B_{1,4}}{B_{2,4} B_{3,4} B_{2,3}} \\
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) s B_{1,2} B_{2,3}}{B_{2,4} B_{3,4} B_{1,4}} - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) B_{2,4} B_{1,2} s}{B_{3,4} B_{2,3} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{2,3} - A_{2,4}) B_{3,4} s B_{1,4}}{B_{2,3} B_{2,4} B_{1,2}} + \frac{1}{64} \frac{(A_{1,4} - A_{2,4}) B_{3,4} s B_{2,3}}{B_{2,4} B_{1,2} B_{1,4}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3}) B_{3,4} B_{2,4} s}{B_{1,2} B_{1,4} B_{2,3}}
\end{aligned}$$

>  $g142334c1324 := \text{map}(\text{factor}, \text{collect}(\text{g142334}\cdot\text{subs}(\text{Sub}, C(1, 3, 2, 4)), [s, B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], \text{distributed}))$

$B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}]$ , distributed)

$$\begin{aligned}
 g142334c1324 := & \frac{1}{64} \frac{(A_{2,3} - A_{3,4})(A_{1,4} - A_{3,4})(A_{1,4} - A_{2,3})(-A_{2,4} + A_{1,3})^2 s^5}{B_{2,3} B_{1,4} B_{3,4} B_{2,4} B_{1,3}} \\
 & - \frac{1}{64} \frac{(A_{2,3} - A_{3,4})(A_{1,4} - A_{3,4})(A_{1,4} - A_{2,3})s^3 B_{1,3}}{B_{2,3} B_{1,4} B_{2,4} B_{3,4}} \\
 & - \frac{1}{64} \frac{(A_{2,3} - A_{3,4})(-A_{2,4} + A_{1,3})^2 s^3 B_{1,4}}{B_{2,4} B_{1,3} B_{2,3} B_{3,4}} + \frac{1}{64} \frac{(A_{1,4} - A_{3,4})(-A_{2,4} + A_{1,3})^2 s^3 B_{2,3}}{B_{2,4} B_{1,3} B_{1,4} B_{3,4}} \\
 & - \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(-A_{2,4} + A_{1,3})^2 B_{3,4} s^3}{B_{2,3} B_{1,4} B_{1,3} B_{2,4}} \\
 & - \frac{1}{64} \frac{(A_{2,3} - A_{3,4})(A_{1,4} - A_{3,4})(A_{1,4} - A_{2,3})B_{2,4} s^3}{B_{3,4} B_{1,4} B_{1,3} B_{2,3}} + \frac{1}{64} \frac{(A_{2,3} - A_{3,4})s B_{1,3} B_{1,4}}{B_{2,4} B_{3,4} B_{2,3}} \\
 & - \frac{1}{64} \frac{(A_{1,4} - A_{3,4})s B_{1,3} B_{2,3}}{B_{2,4} B_{3,4} B_{1,4}} + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})B_{3,4} B_{1,3} s}{B_{2,3} B_{2,4} B_{1,4}} \\
 & + \frac{1}{64} \frac{(A_{2,3} - A_{3,4})B_{2,4} s B_{1,4}}{B_{2,3} B_{3,4} B_{1,3}} - \frac{1}{64} \frac{(A_{1,4} - A_{3,4})B_{2,4} s B_{2,3}}{B_{3,4} B_{1,3} B_{1,4}} \\
 & + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})B_{3,4} B_{2,4} s}{B_{2,3} B_{1,3} B_{1,4}}
 \end{aligned} \tag{40}$$

>  
 $GC0 := m1 \cdot g121324c1423 + m2 \cdot g121334c1423 + m3 \cdot g121423c1324 + m4 \cdot g121434c1324 + m5 \cdot g122334c1324 + m6 \cdot g122434c1423 + m7 \cdot g131423c1234 + m8 \cdot g131424c1234 + m9 \cdot g132324c1234 + m10 \cdot g132434c1423 + m11 \cdot g142324c1234 + m12 \cdot g142334c1324 :$

>  
 $GC := collect(subs(m1=1, m2=1, m3=1, m4=1, m5=1, m6=1, m7=1, m8=1, m9=1, m10=1, m11=1, m12=1, GC0), [B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], distributed) :$

>  
 $Eodd3 := sort(map(factor, collect(Eodd2 - GC, [B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], distributed)), [B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}])$

$$\begin{aligned}
 Eodd3 := & \frac{1}{64} \frac{(A_{2,4} - A_{3,4})(A_{1,4} - A_{2,3})s^2 B_{1,2} B_{1,3}}{B_{1,4} B_{2,3} B_{2,4} B_{3,4}} \\
 & - \frac{1}{64} \frac{(A_{2,3} - A_{3,4})(-A_{2,4} + A_{1,3})s^2 B_{1,2} B_{1,4}}{B_{1,3} B_{2,3} B_{2,4} B_{3,4}} \\
 & + \frac{1}{64} \frac{(A_{1,4} - A_{3,4})(-A_{2,4} + A_{1,3})s^2 B_{1,2} B_{2,3}}{B_{1,3} B_{1,4} B_{2,4} B_{3,4}} \\
 & - \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,3} - A_{3,4})s^2 B_{1,2} B_{2,4}}{B_{1,3} B_{1,4} B_{2,3} B_{3,4}} \\
 & + \frac{1}{64} \frac{(A_{2,3} - A_{2,4})(-A_{3,4} + A_{1,2})s^2 B_{1,3} B_{1,4}}{B_{1,2} B_{2,3} B_{2,4} B_{3,4}} \\
 & - \frac{1}{64} \frac{(A_{1,4} - A_{2,4})(-A_{3,4} + A_{1,2})s^2 B_{1,3} B_{2,3}}{B_{1,2} B_{1,4} B_{2,4} B_{3,4}}
 \end{aligned} \tag{41}$$

$$\begin{aligned}
& + \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{2,4})s^2 B_{1,3} B_{3,4}}{B_{1,2} B_{1,4} B_{2,3} B_{2,4}} \\
& + \frac{1}{64} \frac{(A_{1,3} - A_{2,3})(-A_{3,4} + A_{1,2})s^2 B_{1,4} B_{2,4}}{B_{1,2} B_{1,3} B_{2,3} B_{3,4}} \\
& - \frac{1}{64} \frac{(-A_{2,4} + A_{1,3})(A_{1,2} - A_{2,3})s^2 B_{1,4} B_{3,4}}{B_{1,2} B_{1,3} B_{2,3} B_{2,4}} \\
& - \frac{1}{64} \frac{(A_{1,3} - A_{1,4})(-A_{3,4} + A_{1,2})s^2 B_{2,3} B_{2,4}}{B_{1,2} B_{1,3} B_{1,4} B_{3,4}} \\
& + \frac{1}{64} \frac{(-A_{2,4} + A_{1,3})(A_{1,2} - A_{1,4})s^2 B_{2,3} B_{3,4}}{B_{1,2} B_{1,3} B_{1,4} B_{2,4}} \\
& - \frac{1}{64} \frac{(A_{1,4} - A_{2,3})(A_{1,2} - A_{1,3})s^2 B_{2,4} B_{3,4}}{B_{1,2} B_{1,3} B_{1,4} B_{2,3}} - \frac{1}{64} \frac{1}{B_{1,3} B_{1,4} B_{2,3} B_{2,4} B_{3,4}} ((A_{1,4} \\
& - A_{2,3})(-A_{2,4} + A_{1,3})(-A_{1,3} A_{3,4} + A_{1,3} A_{2,4} + A_{1,4} A_{3,4} - A_{2,3} A_{1,4} - A_{2,4} A_{3,4} \\
& + A_{3,4} A_{2,3})s^4 B_{1,2}) - \frac{1}{64} \frac{1}{B_{1,2} B_{1,4} B_{2,3} B_{2,4} B_{3,4}} ((A_{1,4} - A_{2,3})(-A_{3,4} \\
& + A_{1,2})(A_{1,2} A_{2,4} - A_{1,2} A_{3,4} - A_{1,4} A_{2,4} + A_{2,3} A_{1,4} + A_{2,4} A_{3,4} - A_{2,3} A_{2,4})s^4 B_{1,3}) \\
& + \frac{1}{64} \frac{1}{B_{1,2} B_{1,3} B_{2,3} B_{2,4} B_{3,4}} ((-A_{2,4} + A_{1,3})(-A_{3,4} + A_{1,2})(-A_{1,2} A_{3,4} + A_{1,2} A_{2,3} \\
& + A_{1,3} A_{2,4} - A_{1,3} A_{2,3} + A_{3,4} A_{2,3} - A_{2,3} A_{2,4})s^4 B_{1,4}) - \frac{1}{64} \frac{1}{B_{1,2} B_{1,3} B_{1,4} B_{2,4} B_{3,4}} (( \\
& - A_{2,4} + A_{1,3})(-A_{3,4} + A_{1,2})(-A_{1,2} A_{3,4} + A_{1,2} A_{1,4} + A_{1,3} A_{2,4} - A_{1,3} A_{1,4} + A_{1,4} A_{3,4} \\
& - A_{1,4} A_{2,4})s^4 B_{2,3}) + \frac{1}{64} \frac{1}{B_{1,2} B_{1,3} B_{1,4} B_{2,3} B_{3,4}} ((A_{1,4} - A_{2,3})(-A_{3,4} + A_{1,2})( \\
& - A_{1,2} A_{3,4} + A_{1,3} A_{1,2} + A_{2,3} A_{1,4} - A_{1,3} A_{1,4} + A_{1,3} A_{3,4} - A_{1,3} A_{2,3})s^4 B_{2,4}) \\
& - \frac{1}{64} \frac{1}{B_{1,2} B_{1,3} B_{1,4} B_{2,3} B_{2,4}} ((A_{1,4} - A_{2,3})(-A_{2,4} + A_{1,3})(-A_{1,3} A_{2,4} + A_{1,3} A_{1,2} \\
& + A_{2,3} A_{1,4} - A_{1,2} A_{1,4} + A_{1,2} A_{2,4} - A_{1,2} A_{2,3})s^4 B_{3,4}) \\
& + \frac{1}{64} \frac{1}{B_{1,2} B_{1,3} B_{1,4} B_{2,3} B_{2,4} B_{3,4}} ((A_{1,4} - A_{2,3})(-A_{2,4} + A_{1,3})(-A_{3,4} + A_{1,2})( \\
& - A_{1,3} A_{1,2} A_{3,4} - A_{1,2} A_{1,4} A_{2,3} + A_{1,4} A_{1,2} A_{3,4} - A_{1,2} A_{2,4} A_{3,4} + A_{1,3} A_{1,2} A_{2,4} + A_{1,2} A_{2,3} A_{3,4} \\
& + A_{1,3} A_{1,4} A_{2,3} - A_{1,3} A_{2,3} A_{2,4} + A_{2,3} A_{1,4} A_{2,4} - A_{1,3} A_{1,4} A_{2,4} - A_{1,4} A_{3,4} A_{2,3} \\
& + A_{1,3} A_{2,4} A_{3,4})s^6)
\end{aligned}$$

>  $nops(\%)$

19

(42)

>  $g122434 \cdot g131423$

$$\begin{aligned}
& \left( \frac{1}{32} \frac{(A_{2,4} - A_{3,4}) B_{1,2} s}{B_{3,4} B_{2,4}} - \frac{1}{32} \frac{(-A_{3,4} + A_{1,2}) B_{2,4} s}{B_{3,4} B_{1,2}} + \frac{1}{32} \frac{(A_{1,2} - A_{2,4}) B_{3,4} s}{B_{2,4} B_{1,2}} \right. \\
& \left. - \frac{1}{32} \frac{(A_{2,4} - A_{3,4})(-A_{3,4} + A_{1,2})(A_{1,2} - A_{2,4})s^3}{B_{1,2} B_{2,4} B_{3,4}} \right) \left( \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) B_{1,3} s}{B_{2,3} B_{1,4}} \right.
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{32} \frac{(A_{1,3} - A_{2,3}) B_{1,4} s}{B_{2,3} B_{1,3}} + \frac{1}{32} \frac{(A_{1,3} - A_{1,4}) B_{2,3} s}{B_{1,4} B_{1,3}} \\
& - \frac{1}{32} \frac{(A_{1,4} - A_{2,3}) (A_{1,3} - A_{2,3}) (A_{1,3} - A_{1,4}) s^3}{B_{2,3} B_{1,3} B_{1,4}}
\end{aligned}$$

>  $Eodd4 := sort(map(factor, collect(Eodd3 - 16 \cdot g122434 \cdot g131423 - 16 \cdot g121334 \cdot g142324, [B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}], distributed)), [B_{1,2}, B_{1,3}, B_{1,4}, B_{2,3}, B_{2,4}, B_{3,4}])$

$Eodd4 := 0$  (44)

>  $EvI := \frac{1}{16} \cdot ((1 + C(3, 1, 3, 2) + C(2, 3, 2, 4)) \cdot (1 + C(1, 2, 1, 4) + C(4, 1, 4, 3)) + (1 + C(2, 1, 2, 3) + C(3, 2, 3, 4)) \cdot (1 + C(4, 1, 4, 2) + C(1, 3, 1, 4)) + (1 + C(3, 1, 3, 2) + C(1, 3, 1, 4)) \cdot (1 + C(2, 1, 2, 4) + C(4, 2, 4, 3)) + (1 + C(1, 2, 1, 3) + C(3, 1, 3, 4)) \cdot (1 + C(2, 3, 2, 4) + C(4, 1, 4, 2)) + (1 + C(2, 1, 2, 3) + C(1, 2, 1, 4)) \cdot (1 + C(3, 1, 3, 4) + C(4, 2, 4, 3)) + (1 + C(1, 2, 1, 3) + C(2, 1, 2, 4)) \cdot (1 + C(3, 2, 3, 4) + C(4, 1, 4, 3))) + \frac{1}{8} \cdot (C(1, 4, 2, 3) \cdot C(1, 3, 2, 4) - C(1, 4, 2, 3) \cdot C(1, 2, 3, 4) + C(1, 3, 2, 4) \cdot C(1, 2, 3, 4))$

>  $GC := g121324c1423 + g121334c1423 + g121423c1324 + g121434c1324 + g122334c1324 + g122434c1423 + g131423c1234 + g131424c1234 + g132324c1234 + g132434c1423 + g142324c1234 + g142334c1324 :$

> #####  
#####

## ***INTRINSIC FORMULA FOR THE HYPERBOLIC 4 - ATIYAH DETERMINANT***

>  $DD4 := EvI - \frac{(G123 + G124 + G134 + G234)}{2}$   
 $+ (g121324 + g121334 + g121423 + g121434 + g122334 + g122434 + g131423 + g131424 + g132324 + g132434 + g142324 + g142334)$

$+ GC) + 16 \cdot g122434 \cdot g131423 + 16 \cdot g121334$   
 $\cdot g142324 :$

> #####  
#####

> *indets(DD4)*  
 $\{s, A_{1, 2}, A_{1, 3}, A_{1, 4}, A_{2, 3}, A_{2, 4}, A_{3, 4}, B_{1, 2}, B_{1, 3}, B_{1, 4}, B_{2, 3}, B_{2, 4}, B_{3, 4}\}$  (45)

> *PROOF: factor(DD4 - D4a)*  
0 (46)

### > **IMPORTANT REMARK :**

*We have recently realized that the quadratic part 16*  
 $\cdot g122434 \cdot g131423 + 16 \cdot g121334$   
 $\cdot g142324$  *can be rewritten similarly as a triple cosine*  
*products formula (analogous to the formula*  
**for normalized squared volume of euclidean tetrahedra)**

*In fact it is equal to  $\frac{1}{16}$  times the sum of eight products  $C(i,$   
 $j, i, l) \cdot C(j, k, j, l) \cdot C(k, i, k, l)$  over  $1 \leq j < k < l \leq 4.$*

>  $Y0 := \frac{1}{16} \cdot (C(2, 1, 2, 3) \cdot C(3, 1, 3, 4) \cdot C(1, 4, 2, 4) + C(1, 3, 2, 3) \cdot C(1, 4, 3, 4) \cdot C(2, 1,$   
 $2, 4) + C(1, 2, 1, 3) \cdot C(3, 2, 3, 4) \cdot C(1, 4, 2, 4) + C(1, 3, 2, 3) \cdot C(2, 4, 3, 4) \cdot C(1, 2,$   
 $1, 4) + C(1, 2, 1, 3) \cdot C(2, 3, 2, 4) \cdot C(1, 4, 3, 4) + C(2, 1, 2, 3) \cdot C(2, 4, 3, 4) \cdot C(1, 3,$   
 $1, 4) + C(1, 2, 1, 4) \cdot C(2, 3, 2, 4) \cdot C(3, 1, 3, 4) + C(2, 1, 2, 4) \cdot C(3, 2, 3, 4) \cdot C(1, 3,$   
 $1, 4))$

$$Y0 := \frac{1}{16} \left( \frac{2(t_{2,3} - t_{1,2})(t_{2,1} - t_{3,2})}{(t_{2,1} - t_{1,2})(t_{2,3} - t_{3,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{1,3})(t_{3,1} - t_{4,3})}{(t_{3,1} - t_{1,3})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{2,4} - t_{4,1})(t_{1,4} - t_{4,2})}{(t_{1,4} - t_{4,1})(t_{2,4} - t_{4,2})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{2,3} - t_{3,1})(t_{1,3} - t_{3,2})}{(t_{1,3} - t_{3,1})(t_{2,3} - t_{3,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{4,1})(t_{1,4} - t_{4,3})}{(t_{1,4} - t_{4,1})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{2,4} - t_{1,2})(t_{2,1} - t_{4,2})}{(t_{2,1} - t_{1,2})(t_{2,4} - t_{4,2})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{1,3} - t_{2,1})(t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1})(t_{1,3} - t_{3,1})} - 1 \right) \left( \frac{2(t_{3,4} - t_{2,3})(t_{3,2} - t_{4,3})}{(t_{3,2} - t_{2,3})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{2,4} - t_{4,1})(t_{1,4} - t_{4,2})}{(t_{1,4} - t_{4,1})(t_{2,4} - t_{4,2})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{2,3} - t_{3,1})(t_{1,3} - t_{3,2})}{(t_{1,3} - t_{3,1})(t_{2,3} - t_{3,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{4,2})(t_{2,4} - t_{4,3})}{(t_{2,4} - t_{4,2})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{1,4} - t_{2,1})(t_{1,2} - t_{4,1})}{(t_{1,2} - t_{2,1})(t_{1,4} - t_{4,1})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{1,3} - t_{2,1})(t_{1,2} - t_{3,1})}{(t_{1,2} - t_{2,1})(t_{1,3} - t_{3,1})} - 1 \right) \left( \frac{2(t_{2,4} - t_{3,2})(t_{2,3} - t_{4,2})}{(t_{2,3} - t_{3,2})(t_{2,4} - t_{4,2})} - 1 \right) \left( \frac{2(t_{3,4} - t_{4,1})(t_{1,4} - t_{4,3})}{(t_{1,4} - t_{4,1})(t_{3,4} - t_{4,3})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{2,3} - t_{1,2})(t_{2,1} - t_{3,2})}{(t_{2,1} - t_{1,2})(t_{2,3} - t_{3,2})} - 1 \right)$$
 (47)

$$\begin{aligned}
& -1 \left( \frac{2(t_{3,4} - t_{4,2})(t_{2,4} - t_{4,3})}{(t_{2,4} - t_{4,2})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{1,4} - t_{3,1})(t_{1,3} - t_{4,1})}{(t_{1,3} - t_{3,1})(t_{1,4} - t_{4,1})} - 1 \right) \\
& + \frac{1}{16} \left( \frac{2(t_{1,4} - t_{2,1})(t_{1,2} - t_{4,1})}{(t_{1,2} - t_{2,1})(t_{1,4} - t_{4,1})} - 1 \right) \left( \frac{2(t_{2,4} - t_{3,2})(t_{2,3} - t_{4,2})}{(t_{2,3} - t_{3,2})(t_{2,4} - t_{4,2})} \right. \\
& \left. - 1 \right) \left( \frac{2(t_{3,4} - t_{1,3})(t_{3,1} - t_{4,3})}{(t_{3,1} - t_{1,3})(t_{3,4} - t_{4,3})} - 1 \right) + \frac{1}{16} \left( \frac{2(t_{2,4} - t_{1,2})(t_{2,1} - t_{4,2})}{(t_{2,1} - t_{1,2})(t_{2,4} - t_{4,2})} \right. \\
& \left. - 1 \right) \left( \frac{2(t_{3,4} - t_{2,3})(t_{3,2} - t_{4,3})}{(t_{3,2} - t_{2,3})(t_{3,4} - t_{4,3})} - 1 \right) \left( \frac{2(t_{1,4} - t_{3,1})(t_{1,3} - t_{4,1})}{(t_{1,3} - t_{3,1})(t_{1,4} - t_{4,1})} - 1 \right)
\end{aligned}$$

1

>  $Y1 := \text{sort}\left(\text{map}\left(\text{factor}, \text{collect}\left(\text{map}\left(\text{factor}, \text{collect}\left(\text{factor}\left(\text{expand}\left(\text{subs}\left(\text{seq}\left(\text{seq}\left(t[i, j] = \frac{(s \cdot A[i, j] + B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right), \text{seq}\left(\text{seq}\left(t[j, i] = \frac{(s \cdot A[i, j] - B[i, j])}{2}, i = 1 .. j - 1\right), j = 1 .. 6\right)\right), Y0\right)\right), s\right)\right), [B_{1, 2}, B_{1, 3}, B_{1, 4}, B_{2, 3}, B_{2, 4}, B_{3, 4}], \text{distributed}\right)\right), s\right) :$

> *factor*(*Eodd3* - *Y1*)

0

(48)

2

(49)