SPECIAL *n***th ORDER SURFACES WITH** (n-2)-**ple LINE**

Sonja GORJANC

University of Zagreb, Croatia

ABSTRACT: In this paper, in Euclidean space \mathbb{E}^3 , we treat the pedal surfaces of special line congruences C_{2k}^1 which are of the 1st order and the 2kth class. We derive the parametric and implicit equations of these surfaces which enable *Mathematica* visualizations and proving some properties such as their order is 2k + 2, they possess one 2k-ple straight line and pass through the absolute conic of \mathbb{E}^3 . The properties of their singularities, which do not lie on 2k-ple line, and of the pinch points on the 2k-ple line, are also shown.

Keywords: congruence of lines, inversion, pedal surfaces of congruence, multiple line, multiple point, pinch point

1. INTRODUCTION

Congruence C is a double infinite line system, i.e. it is the set of lines in the threedimensional space (projective, affine or Euclidean) depending on two parameters. Line $z \in C$ is said to be the *ray* of a congruence. The *order* of a congruence is the number of its rays which pass through an arbitrary point; the *class* of a congruence is the number of its rays which lie in an arbitrary plane. *mth order*, *nth class* congruence is signed C_n^m . A point is called the *singular point* of a congruence if ∞^1 rays pass through it. A plane is called the *singular plane* of a congruence if it contains ∞^1 rays.

According to [6, p. 64], [10, pp. 1184-1185], there are only two types of the first order congruences: the first one are the congruences of nth class and their rays are transversals of one straight line d and nth order space curve c^n



Figure 1: Directing lines of C_n^1

which cuts this straight line in n - 1 points (see Fig. 1), and the second are congruences of 3rd class and its rays cut a twisted cubic twice. The properties of the first order congruences can be found in [1].

In Euclidean space E^3 , the *pedal surface* of congruence C_n^m with respect to *pole* P is the locus of the feet of perpendiculars from finite point P to the rays of congruence C_n^m , [5].

In [2] we define the transformation of threedimensional projective space where corresponding points lie on the rays of congruence C_n^1 and are conjugate with respect to proper quadric Ψ (see Fig. 2). This transformation we called the (n + 2) degree inversion with respect to congruence C_n^1 and quadric Ψ and signed it by $i_{\Psi}^{n+2} : \mathbb{P}^3 \to \mathbb{P}^3$. We proved that it takes a straight line to the (n + 2) order space



Figure 2: Inversion of degree n + 2

curve and a plane to the (n+2) order surface which contains *n*-ple straight line.

The class of such surfaces was elaborated in detail by Sturm [8, pp. 315-328].

The pedal surface of the first order congruence C_n^1 is the image of the plane at infinity given by i_{Ψ}^{n+2} , where Ψ is any sphere with center P. According to the properties of i_{Ψ}^{n+2} it was shown that the pedal surface of congruence C_n^1 is (n+2) order surface with *n*-ple line straight line *d* which passes through the absolute conic of E^3 and the directing curve c^n , [2].

2. SPECIAL C_{2k}^1 **CONGRUENCES**

A special class of C_n^1 arises if all intersection points D_i (see Fig. 1) coincide. In this case c^n is a plane curve with one singular point of the highest multiplicity n - 1, and line d passes through this point.

Here we will regard special C_n^1 where n is an even number, i. e. $n = 2k, k \in \mathbb{N}$, and directing curve c^{2k} is a plane curve with (2k-1)-ple singular point.

2.1 (2k - 1)-folium

(2k-1)-folium is curve c^{2k} given by the following polar equation:



Figure 3: (2k - 1)-folium

According to the multiple-angle formula, $\cos(2k-1)\varphi$ can be displayed as

$$\sum_{i=0}^{k} (-1)^{i} C_{2i}^{2k-1} (\cos \varphi)^{2k-1-2i} (\sin \varphi)^{2i} \quad (2)$$

where C_{2i}^{2k-1} is a binomial coefficient.

Therefore, from eq. (1), by using the substitutions $r(\varphi) = \sqrt{x^2 + y^2}$, $\cos \varphi = \frac{x}{\sqrt{x^2 + y^2}}$ and $\sin \varphi = \frac{y}{\sqrt{x^2 + y^2}}$, we obtain the following implicit equation of (2k - 1)-folium:

$$(x^2 + y^2)^k - \tau^{2k-1} = 0$$
, where, (3)

$$\tau^{2k-1} = \sum_{i=0}^{k} (-1)^{i} C_{2i}^{2k-1} x^{2k-1-2i} y^{2i}.$$
 (4)

From eq. (3) it is clear that (2k - 1)-folium is 2k-order curve c^{2k} , with (2k - 1)-ple point at the origin, where 2k - 1 tangent lines at it are given by equation $\tau^{2k-1} = 0$, [7, p. 27]. The line at infinity is the k-ple tangent line of c^{2k} which touches it at the absolute points.

2.2 Congruence C_{2k}^1

Let axis z and (2k - 1)-folium c^{2k} in plane z = 0 be the directing lines of congruence C_{2k}^1 .



Figure 4: Directing lines of C_{2k}^1 for k = 2

All singular points of C_{2k}^1 (the points which contain ∞^1 rays of C_{2k}^1) lie on its directing lines c^{2k} and z. If point C lies on curve c^{2k} and $C \neq O$, then the rays of C_{2k}^1 which pass trough C form pencil of lines (C) in plane $\zeta \in [z]$ which contains C and z, see Fig. 5a. If point Z lies on axis z and $Z \neq O$, then all the lines which join Z with the points of curve c^{2k} are



Figure 5: Singular points and planes of C_{2k}^1 for k = 2.



Figure 6: Rays of C_{2k}^1 (for k = 2) through non singular points and non singular planes.

the rays of C_n^1 . They form 2k- degree cone Φ_Z^{2k} with vertex Z. Axis z is (2k - 1)-ple generatrix of Φ_Z^{2k} , see Fig. 5b. The rays through point O form 2k - 1 pencil of lines (O) in the planes determined by axis z and 2k - 1 tangent lines of c^{2k} at O, and pencil (O) in the plane of c^{2k} , see Fig. 5c. Singular planes of C_{2k}^1 (the planes which contain ∞^1 rays) are the planes of the pencil [z] and plane of (2k - 1)-folium, see Fig. 5c.

Every point A, which is not the singular point of C_{2k}^1 , determines plane $\zeta_A \in [z]$ which cuts c^{2k} in only one point C beside the origin. Line AC, which cuts z in one point Z, is the unique ray of C_{2k}^1 through point A, see Fig. 6a. If plane ζ_A contains one of the tangent lines of c^{2k} at O, then points C and Z coincide with O and line AO is the unique ray of C_{2k}^1 through A, see Fig. 6b.

Every plane α which is not the singular plane of C_{2k}^2 , contains 2k rays of C_{2k}^1 . Plane α cuts axis z in point Z and curve c^{2k} in points $C_j, j = 1, ..., 2k$. Lines ZC_j are 2k rays of C_n^1 in plane α . They are the intersection of plane α and 2k-degree cone Φ_Z^{2k} , and can be real and different, coinciding or imaginary, see Fig. 6c.. If α passes through point O, then 2k - 1 rays are the intersections of α with the planes through z and the tangent lines of c^{2k} at O, and one ray lies in the plane of c^{2k} , see Fig. 6d.

3. PEDAL SURFACES OF C_{2k}^1

As we mentioned in the Introduction: in Euclidean space \mathbb{E}^3 , the pedal surface of congruence \mathcal{C} with respect to pole P is the locus of the feet of perpendiculars from finite point P to the rays of congruence \mathcal{C} . According to [2], the pedal surfaces of \mathcal{C}_{2k}^1 is 2k + 2 order surface with 2k-ple line z, and we will denote it \mathcal{P}_{2k}^{2k+2} .



Figure 7: Construction of circles $c \in \zeta \in [z]$

3.1 Construction

In plane ζ trough axis z, the rays of C_{2k}^1 form pencil of lines (C), where $C \neq O$ is the intersection of ζ and c^{2k} . In 2k - 1 planes, determined by the tangent lines of c^{2k} at O and axis z, point C coincides with O. If finite pole P is in the general position to the directing lines of C_{2k}^1 , the feet of perpendiculars from P to the rays of pencil (C) form circle c with diameter $\overline{CP'}$, where P' is the orthogonal projection of P to ζ , see Fig. 7a,b. The proof of this statement is elementary.

For given pole P, the path of point P', with respect to the moving plane ζ , is the circle, denoted by k, which lies in the plane through P, perpendicular to axis z. The diameter of kis $\overline{PP_z}$, where P_z is the normal projection of P to z. Thus, we can regard surface \mathcal{P}_{2k}^{2k+2} as the system of circles in the planes through axis z with the end points of diameters on (2k-1)folium c^{2k} and circle k, see Fig. 7c,d.

The diameters of circles c lie on ruled surface with directing lines c^{2k} , k and z. According to the formula [6, p. 90], the degree of this surface is: $2 \cdot 2k \cdot 2 \cdot 1 - 2 \cdot 1 - (2k-1) \cdot 2 - 1 \cdot 2k = 2k$.

3.2 Parametric equations of \mathcal{P}_{2k}^{2k+2} and *Mathematica* visualizations

Let $(p_x, p_y, p_z) \in \mathbb{R}^3$ be the coordinates of pole P and let (2k - 1)-folium is given by eq. (1). Let (r, z), where $|r| = \sqrt{x^2 + y^2}$, be the coordinates of the points in plane $\zeta(\varphi)$, which is given by equation $y = x \tan \varphi$ if $\varphi \in [0, \pi), \varphi \neq \pi/2$, and x = 0 if $\varphi = \pi/2$, see Fig. 8.



Figure 8

The coordinates of points $P', C \in \zeta(\varphi)$ are

$$(r_{P'}, z_{P'})(\varphi) = (p_x \cos \varphi + p_y \sin \varphi, p_z) (r_C, z_C)(\varphi) = (\cos(2k - 1)\varphi, 0).$$
 (5)

 $R(\varphi)$ is the radius and $S(r_S(\varphi), z_S(\varphi))$ is the center of circle c in plane $\zeta(\varphi)$.

$$R(\varphi) = \frac{1}{2}\sqrt{(r_C(\varphi) - r_{P'}(\varphi))^2 + p_z^2}$$

$$r_S(\varphi) = \frac{r_C(\varphi) + r_{P'}(\varphi)}{2}$$

$$z_S(\varphi) = \frac{p_z}{2}.$$
(6)

Since the parametric equations of circle c in plane $\zeta(\varphi)$ are

$$r(\theta) = R(\varphi) \sin \theta + r_S(\varphi)$$

$$z(\theta) = R(\varphi) \cos \theta + z_S(\varphi),$$

$$\theta \in [0, 2\pi),$$
(7)



Figure 9: \mathcal{P}_{2k}^{2k+2} for P(1,0,2) and k = 1, 2, 3, 4, respectively in figures a, b, c and d

therefore the parametric equations of surface \mathcal{P}^{2k+1}_{2k} are the following

$$\begin{aligned} x(\varphi,\theta) &= \cos\varphi \left(R(\varphi)\sin\theta + r_S(\varphi) \right) \\ y(\varphi,\theta) &= \sin\varphi \left(R(\varphi)\sin\theta + r_S(\varphi) \right) \\ z(\varphi,\theta) &= R(\varphi)\cos\theta + z_S(\varphi), \\ \varphi \in [0,\pi), \ \theta \in [0,2\pi). \end{aligned}$$

Equations (8) enable *Mathematica* visualizations of surfaces \mathcal{P}_{2k}^{2k+1} . See Fig. 9.

3.3 Implicit equation of \mathcal{P}^{2k+2}_{2k}

In plane $\zeta(u)$ through axis z, in coordinates (r, z), the equation of circle c is

$$(r - r_S(\varphi))^2 + (z - p_z/2)^2 = R(\varphi)^2,$$

 $\varphi \in [0, \pi).$ (9)

From equations (5), by using the formula (2) and the substitutions $\cos u = \frac{x}{\sqrt{x^2+y^2}}$, $\sin u = \frac{y}{\sqrt{x^2+y^2}}$, we obtain the following

$$r_{C}(\varphi) = \frac{\tau^{2k-1}}{\sqrt{(x^{2}+y^{2})^{2k-1}}}$$
$$r_{P'}(\varphi) = \frac{p \, x + q y}{\sqrt{x^{2}+y^{2}}} \tag{10}$$

where τ^{2k-1} is given by eq. (4).

Now, we can express $r_S(\varphi)$ and $R(\varphi)$, given by formulas (6), as the functions of x and y. If we put these functions and $r = \sqrt{x^2 + y^2}$ into equation (9) and multiply it by $(x^2 + y^2)^k$, we obtain the implicit equation of \mathcal{P}^{2k+2}_{2k} which can be written in the following form

$$(x^{2} + y^{2})^{k}(x^{2} + y^{2} + z^{2}) + H^{2k+1}(x, y) + H^{2k}_{1}(x, y)z + H^{2k}_{2}(x, y) = 0,$$
(11)

where $H^i(x, y)$ are homogeneous polynomials in x and y of degree i, given by the formulas:

$$H^{2k+1}(x,y) = - (x^{2} + y^{2})^{k} (p_{x}x + p_{y}y) - (x^{2} + y^{2})\tau^{2k-1} H^{2k}_{1}(x,y) = -p_{z}(x^{2} + y^{2})^{k} H^{2k}_{2}(x,y) = (p_{x}x + p_{y}y)\tau^{2k-1}.$$
 (12)

3.4 Properties of \mathcal{P}^{2k+2}_{2k}

Proposition 1 The plane at infinity cuts surface \mathcal{P}_{2k}^{2k+2} at the absolute conic of \mathbb{E}^3 and the rays of congruence \mathcal{C}_{2k}^1 .

PROOF: In the Cartesian homogeneous coordinates (x : y : z : w), where w = 0 means that the point lies in the plane at infinity, the equation of surface \mathcal{P}_{2k}^{2k+2} takes the form

$$(x^{2} + y^{2})^{y}(x^{2} + y^{2} + z^{2}) + H^{2k+1}(x, y)w + H^{2k}_{1}(x, y)zw + H^{2k}_{2}(x, y)w^{2} = 0.$$
(13)

Therefore, the intersection of \mathcal{P}_{2k}^{2k+2} and the plane at infinity splits into the absolute conic, given by equations $x^2 + y^2 + z^2 = 0$, w = 0, and the pair of imaginary lines through the

point (0:0:1:0), counted k times, which are given by equations $(x^2 + y^2)^k = 0$, w = 0. It is clear from eq. (3) that curve c^{2k} touches the plane at infinity k times at the absolute points. Thus, C_{2k}^1 has k-ple pair of isotropic rays through (0:0:1:0) at infinity. \Box

Proposition 2 Axis z is the 2k-ple line of surface \mathcal{P}_{2k}^{2k+2} .

PROOF: According to [4, p. 251]: If the *n*th order surface in \mathbb{E}^3 , which passes through the origin, is given by equation

 $F(x, z, y) = f_m(x, y, z) + f_{m+1}(x, y, z) + \cdots$ $\cdots + f_n(x, y, z) = 0$, where $f_k(x, y, z)$ $(1 \le k \le n)$ is homogeneous polynomial of degree k, then the tangent cone at the point (0, 0, 0) is given by equation $f_m(x, y, z) = 0$.

If we move the origin to any point $Z_0 = (0, 0, z_0)$ on axis z, from eq. (11) we obtain the following equation for the tangent cone \mathcal{T}_{Z_0} of \mathcal{P}_{2k}^{2k+2} at point Z_0

$$(x^{2} + y^{2})^{k} z_{0}^{2} + H_{1}^{2k}(x, y) z_{0} + H_{2}^{2k}(x, y) = 0.$$
(14)

Since it is the homogeneous equation in x and y of degree 2k, in the general case \mathcal{T}_{Z_0} always splits into 2k planes through axis z.

There are many possibilities for the type of 2kple singular point Z_0 on line z. It depends on how the homogeneous polynomial from eq. (14) can be factorized, i. e. how tangent cone \mathcal{T}_{Z_0} splits (how many real and imaginary planes, how many coinciding planes, and so on). For example, for point O, \mathcal{T}_O is given by equation $(p_x x + p_y y)\tau^{2k-1} = 0$ and, in general, splits into 2k real and different planes. But, if the line in plane xy which is given by $p_x x + p_y y = 0$ coincides with one of the tangent lines of c^{2k} through O, \mathcal{P}_{2k}^{2k+2} has the *pinch point* in O. Pinch points are the points on multiple line in which two or more tangent planes coincide.

Proposition 3 Surface \mathcal{P}_{2k}^{2k+2} has 4(2k-1) pinch points on 2k-ple axis z (real or complex). Among them one is always the point at infinity and it is the pinch-point counted k times.

PROOF: The proof that *n*th order surface with (n-2)-ple line always possesses 4(n-3)pinch-points, is given in [8, p. 317]. We give here only its interpretation for this 2korder case: Every plane ζ through axes z cuts \mathcal{P}^{2k+2}_{2k} into the 2k-ple line and one conic cwhich cuts 2k-ple line in two points. These points are the touching points of plane ζ and surface \mathcal{P}_{2k}^{2k+2} . The correspondence between the planes of pencil [z], where corresponding planes have the same touching point, is the involution of the order 2(2k-1), because that through each touching point of plane ζ another 2k-1 tangent planes pass. This involution has $2 \cdot 2(2k-1)$ double elements which are the coinciding tangent planes through the points on 2k-ple line and their touching points are the pinch-points of \mathcal{P}^{2k+2}_{2k} .

According to eq. (13), the tangent cone at point $Z_0^{\infty}(0:0:1:0)$ is given by equation $(x^2 + y^2)^k = 0$, thus Z_0^{∞} is the pinch-point counted k times.

Proposition 4 \mathcal{P}_{2k}^{2k+2} contains curve c^{2k} .

PROOF: If z = 0. eq. (11) takes the form

$$(x^{2}+y^{2}-p_{x}x-p_{y}y)((x^{2}+y^{2})^{k}-\tau^{2k+1})=0.$$

Thus, plane z = 0 cuts \mathcal{P}_{2k}^{2k+2} through curve c^{2k} and circle with diameter OP', where P' is the normal projection of pole P on plane z = 0.

Proposition 5 If pole P lies on axis z, \mathcal{P}_{2k}^{2k+2} splits into the pair of isotropic planes through z and 2k-order surface.

PROOF: If $p_x = p_y = 0$, eq. (11) takes the form

$$(x^2 + y^2)P^{2k}(x, y, z) = 0,$$

where $P^{2k}(x, y, z)$ is $(x^2 + y^2)^{k-1}(x^2 + y^2 + z^2 - p_z z) - \tau^{2k-1}$. \Box See figures 10 and 11.



Figure 10: Surfaces \mathcal{P}^{2k} , given by equations $P^{2k}(x, y, z) = 0$, for k = 2, 3, 4 and P(0, 0, 0)



Figure 11: Surfaces \mathcal{P}^{2k} , given by equations $P^{2k}(x, y, z) = 0$, for k = 2, 3, 4 and P(0, 0, 2)

Proposition 6 Surface \mathcal{P}_{2k}^{2k+2} has real double points out of axis z iff $p_z = 0$. There are at the most 2k - 1 and at least 1 such real points on \mathcal{P}_{2k}^{2k+2} .

PROOF:

Except the points of 2k-ple line z, the highest singularity which \mathcal{P}_{2k}^{2k+2} can possess is a double point. Namely, if \mathcal{P}_{2k}^{2k+2} had a higher multiple point out of z, the line through that point which cuts z would cut \mathcal{P}_{2k}^{2k+2} in more than 2k + 2 points, which is impossible.

If D is the double point of \mathcal{P}_{2k}^{2k+2} it is the double point of every section of \mathcal{P}_{2k}^{2k+2} through D. Thus, circle c in the plane ζ through D and axis z splits into the pair of isotropic lines through D. It is the case when the end points of diameter $\overline{CP'}$ coincide, i. e. circle k intersects curve c^{2k} . If $p_z \neq 0$ circle k and curve c^{2k} intersect only into the absolute points of

plane z = 0. Curves k and c^{2k} can possess real intersection points only in the case when $p_z = 0$. In this case they have 4k intersection points, where 2k - 1 points coincide with O, 2points are the absolute points of plane z = 0, thus only 2k - 1 intersection points can lie out of axis z and be real. Since 2k - 1 is an odd number, at least one real double point exists on \mathcal{P}_{2k}^{2k+2} , if $p_z=0$.

See figures 12 and 13.



Figure 12: Surface \mathcal{P}_2^4 with 1 real double point out of its double line.



Figure 13: Surfaces \mathcal{P}_4^6 with 1, 2 and 3 real double points out of their quadruple lines

4. CONCLUSIONS

The objective of this paper is to visualize numerous forms and properties of special class of surfaces in \mathbb{E}^3 . Surfaces treated in this paper form only a small subclass of \mathbb{P}_n^{n+2} which is the class of the (n+2)th surfaces with *n*-ple straight line. It may be assumed that the whole class \mathbb{P}_n^{n+2} could be obtained by inversion i_{Ψ}^{n+2} and visualized by the program *Mathematica*..

ACKNOWLEDGMENTS

I would like to thank my colleagues Vladimir Benić and Miklós Hoffmann for their generous support and constructive corrections.

REFERENCES

- [1] V. Benić, S. Gorjanc, (1,n) Congruences, *KoG*, *10* (2006), 5-12.
- [2] V. Benić, S. Gorjanc, Inversion of Degree n + 2, (manuscript - submitted to *Acta Mathematica Hungarica*)
- [3] A. Gray, Modern Differential Geometry of Curves and Surfaces with Mathematica, CRC Press, Boca Raton, 1998.
- [4] J. Harris, *Algebraic Geometry*. Springer, New York, 1995.
- [5] E. Kranjčević, Die Fusspunktflächen der linearen Kongruenzen, *Glasnik matematički*, 3 (23), (1968), 269-274.

- [6] G. Salmon, A Treatise on the Analytic Geometry of Three Dimensions, Vol.II, Chelsea Publishing Company, New York, 1965.
- [7] G. Salmon, *Higher Plane Curves*, Chelsea Publishing Company, New York, 1960.
- [8] R. Sturm, Die Lehre von den geometrischen Verwandtschaften, Band IV, B. G. Taubner, Leipzig-Berlin, 1909.
- [9] R. Sturm, *Liniengeometrie*, *II. Teil*, B. G. Taubner, Leipzig, 1893.
- [10] K. Zindler, Algebraische Liniengeometrie, Encyklopädie der Mathematischen Wissenschaften, Band III, 2. Teil, 2. Hälfte. A., pp. 1184-1185, B. G. Teubner, Liepzig, 1921-1928.

ABOUT THE AUTHOR

Sonja Gorjanc, PH.D. math., is an assistant professor in the Department of Mathematics, Faculty of Civil Engineering, University of Zagreb and actual president of the Croatian Society for Geometry and Graphics. Her research interest are in Projective and Euclidean geometry, *Mathematica* computer graphics, Curricular Developments in Geometry and Graphics. She can be reached by e-mail: sgorjanc@grad.hr or through postal address: Faculty of Civil Engineering, Kačićeva 26, 10000 Zagreb, Croatia.